

BEST APPROXIMATION FOR CONVEX SUBSETS OF 2-INNER PRODUCT SPACES

M. ABRISHASMI-MOGHADDAM^{1,*} AND T. SISTANI²

¹Department of Mathematics, Birjand Branch, Islamic Azad University, Birjand, Iran

²Department of Mathematics, Kerman Branch, Islamic Azad University, Kerman, Iran

ABSTRACT. In this paper, we study the concept of best approximation in 2-inner product spaces. We get some characteristic theorems for the elements of best approximation for convex subsets of 2-inner product spaces. Finally we get some properties of the metric projection map in this spaces.

KEYWORDS : 2-Inner product space; 2-Normed space; b-Best approximation; b-Proximinal; b-Chebyshev; b-Metric projection; b-Dual cone

AMS Subject Classification: 41A65 41A15

1. INTRODUCTION

Recently, some results on best approximation theory in linear 2-normed spaces have been obtained by Y. J. Cho, S. Elumalai, S. S. Kim, R. Ravi, Sh. Rezapour and others (see [1], [4], [6], [12], [13], [15]). These papers are based on the research works in normed linear spaces made by I. Singer ([14]), T. D. Narang ([11]), S. S. Dragomir ([2]) and others. In this paper we want to investigate the concept of best approximation in 2-inner product spaces. The concept of 2-inner product spaces has been investigated by R. Ehret in 1969([3]), and has been developed extensively in different subjects by others.([10], [8])

Definition 1.1. Let X be a linear space of dimension greater than 1 over field \mathbb{R} of real numbers.

Suppose that $\langle \cdot, \cdot | \cdot \rangle$ is a \mathbb{R} -valued function defined on $X \times X \times X$ satisfying the following conditions:

- $\langle x, x | z \rangle \geq 0$ and $\langle x, x | z \rangle = 0$ if and only if x and z are linearly dependent.
- $\langle x, x | z \rangle = \langle z, z | x \rangle$
- $\langle y, x | z \rangle = \langle x, y | z \rangle$
- $\langle \alpha x, y | z \rangle = \alpha \langle x, y | z \rangle$ for any scalar $\alpha \in \mathbb{R}$
- $\langle x + x', y | z \rangle = \langle x, y | z \rangle + \langle x', y | z \rangle$

* Corresponding author.

Email address : m.abrishami.m@gmail.com(M. Abrishasmi-Moghaddam), Taherehsistani@yahoo.com(T. Sistani).

Article history : Received 14 May 2012. Accepted 28 August 2012.

$\langle \cdot, \cdot | \cdot \rangle$ is called a *2-inner product* and $(X, \langle \cdot, \cdot | \cdot \rangle)$ is called a *2-inner product space* (or a *2-per-Hilbert space*). A concept which is closely related to 2-inner product space and introduced by Gähler in 1965, is 2-normed space [5].

Definition 1.2. Let X be a linear space of dimension greater than 1 over field \mathbb{R} of real numbers. Suppose $\| \cdot, \cdot \|$ is a real-valued function on $X \times X$ satisfying the following conditions:

- $\|x, y\| = 0$ if and only if x and y are linearly dependent vectors.
- $\|x, y\| = \|y, x\|$ for all $x, y \in X$.
- $\|\lambda x, y\| = |\lambda| \|x, y\|$ for all $\lambda \in \mathbb{R}$ and $x, y \in X$.
- $\|x + y, z\| \leq \|x, z\| + \|y, z\|$ for all $x, y, z \in X$.

Then $\| \cdot, \cdot \|$ is called a *2-norm* on X and $(X, \| \cdot, \cdot \|)$ is called a *linear 2-normed space*. It is easy to show that the 2-norm $\| \cdot, \cdot \|$ is non-negative and $\|x, y + \alpha x\| = \|x, y\|$ for all $x, y \in X$ and $\alpha \in \mathbb{R}$. Every 2-normed space is a locally convex topological vector space. In fact for a fixed $b \in X$, $p_b(x) = \|x, b\|$; $x \in X$ is a semi-norm on X and the family $P = \{p_b : b \in X\}$ of seminorms generates a locally convex topology on X .

Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space, then

- We can define a 2-norm on $X \times X$ by $\|x, y\| = \sqrt{\langle x, x | y \rangle}$.
- Let $0 \neq b \in X$ and $x, y \in X \setminus \langle b \rangle$. An element $x \in X$ is said to be *b-orthogonal* to an element $y \in X$, and we write $x \perp_b y$, if $\langle x, y | b \rangle = 0$.
- For all $x, y, b \in X$, we have the Cauchy-Schwartz inequality

$$\langle x, y | b \rangle^2 \leq \|x, b\|^2 \|y, b\|^2.$$

Let $(X, \| \cdot, \cdot \|)$ be a 2-normed space and V_1 and V_2 be two linear subspaces of X . A 2-functional $f : V_1 \times V_2 \rightarrow \mathbb{R}$ is called a *bilinear 2-functional* on $V_1 \times V_2$, whenever for all $x_1, x_2 \in V_1, y_1, y_2 \in V_2$ and $\lambda_1, \lambda_2 \in \mathbb{R}$;

- $f(x_1 + x_2, y_1 + y_2) = f(x_1, y_1) + f(x_1, y_2) + f(x_2, y_1) + f(x_2, y_2)$,
- $f(\lambda_1 x_1, \lambda_2 y_1) = \lambda_1 \lambda_2 f(x_1, y_1)$.

A bilinear 2-functional $f : V_1 \times V_2 \rightarrow \mathbb{R}$ is said to be bounded if there exists a non-negative real number M (called a Lipschitz constant for f) such that $|f(x, y)| \leq M \|x, y\|$ for all $x \in V_1$ and $y \in V_2$. Also, the norm of a bilinear 2-functional f is defined by

$$\|f\| = \inf\{M \geq 0 : M \text{ is a Lipschitz constant for } f\}$$

It is known that

$$\begin{aligned} \|f\| &= \sup\{|f(x, y)| : (x, y) \in V_1 \times V_2, \|x, y\| \leq 1\} \\ &= \sup\{|f(x, y)| : (x, y) \in V_1 \times V_2, \|x, y\| = 1\} \\ &= \sup\{|f(x, y)| / \|x, y\| : (x, y) \in V_1 \times V_2, \|x, y\| > 0\}. \end{aligned}$$

Definition 1.3. [7] A 2-functional $F : V_1 \times V_2 \rightarrow \mathbb{R}$ is said to be a convex 2-functional if

$$\begin{aligned} F(a\lambda x + (a - a\lambda)x', b\mu y + (b - b\mu)y') &\leq ab|\lambda\mu|F(x, y) + a|\lambda|(b - b\mu)F(x, y') \\ &\quad + (a - a\lambda)b|\mu|F(x', y) + (a - a\lambda)(b - b\mu)F(x', y') \end{aligned}$$

for all $|\lambda| \leq 1, |\mu| \leq 1$ and $a, b \geq 0$.

Definition 1.4. Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space, and $b \in X$.

1) A sequence $\{x_n\}$ of X is said to be *b-convergent* and denote by $x_n \xrightarrow{b} x$, if there exists an element $x \in X$ such that $\lim_{n \rightarrow \infty} \|x_n - x, b\| = 0$ for all $x \in X$.

2) A subset E of X is said *b-closed*, if for each sequence $\{x_n\}$ in E such that $x_n \xrightarrow{b} x$, we have that $x \in E$.

Definition 1.5. [13] Let $(X, \|\cdot, \cdot\|)$ be a 2-normed space, G a nonempty subset of X , $0 \neq b \in X$, then $g_0 \in G$ is called a *b-best approximation* to x from G if

$$\|x - g_0, b\| = \inf\{\|x - g, b\| : g \in G\}.$$

The set of all b-best approximations of x in G is denoted by $P_{G,b}(x)$. The mapping $P_{G,b} : X \rightarrow 2^G$ is called the *b-metric projection* onto G .

If each $x \in X \setminus (G + \langle b \rangle)$ has at least (resp. exactly) one b-best approximation in G , then G is called a *b-proximinal* (resp. *b-chebyshev*) set.

Definition 1.6. 1) A nonempty subset K of the 2-inner product space X , is called convex if $\lambda x + (1 - \lambda)y \in K$ whenever $x, y \in K$ and $0 \leq \lambda \leq 1$.

2) A nonempty subset C of the 2-inner product space X , is called a convex cone if $\alpha x + \beta y \in C$ whenever $x, y \in C$ and $0 \leq \alpha, \beta \in \mathbb{R}$.

3) A nonempty subset M of the 2-inner product space X , is called a linear subspace if $\alpha x + \beta y \in M$ whenever $x, y \in M$ and $\alpha, \beta \in \mathbb{R}$.

4) A nonempty subset V of the 2-inner product space X , is called affine if $\alpha x + (1 - \alpha)y \in V$ whenever $x, y \in V$, and $\alpha \in \mathbb{R}$.

Definition 1.7. Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space over the real number field \mathbb{R} . The *b-dual cone* (or *b-negative polar*) of S is the set

$$S_b^\circ := \{x \in X | \langle x, y | b \rangle \leq 0 \text{ for all } y \in S\}$$

The *b-orthogonal complement* of S is the set

$$S_b^\perp = S_b^\circ \cap (-S_b^\circ) = \{x \in X | \langle x, y | b \rangle = 0 \text{ for all } y \in S\}$$

2. CHARACTERIZATION THEOREMS FOR ELEMENTS OF B-BEST APPROXIMATION FOR CONVEX SUBSETS OF A 2-INNER PRODUCT SPACE

In this section we investigate some characteristic theorems for elements of b-best approximation for convex subsets of a 2-inner product space X . It is known that every nonempty b-closed convex set in a b-Hilbert space is b-chebyshev (see [9], [15]). Now we have the following theorem.

Theorem 2.1. Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space, and K a b-closed convex subset of X with $X \neq K$. If $x_0 \in X \setminus (K + \langle b \rangle)$ and $g_0 \in K$, then the following statements are equivalent.

(1) $g_0 = P_{K,b}(x_0)$

(2) $x_0 - g_0 \in (K - g_0)_b^\circ$

(3) we have that

$$\inf_{g \in K} \langle g - x_0, g_0 - x_0 | b \rangle = \|g_0 - x_0, b\|^2$$

Proof. (1) \Rightarrow (2) : This implication is proved in ([15]).

(2) \Rightarrow (3) : By (2) we have that for all $g \in K$,

$$\begin{aligned} 0 &\leq \langle x_0 - g_0, g_0 - g | b \rangle = \langle x_0 - g_0, g_0 - x_0 + x_0 - g | b \rangle \\ &= \langle x_0 - g_0, x_0 - g | b \rangle - \langle x_0 - g_0, x_0 - g_0 | b \rangle \\ &= \langle x_0 - g_0, x_0 - g | b \rangle - \|x_0 - g_0, b\|^2. \end{aligned}$$

Hence

$$\|x_0 - g_0, b\|^2 \leq \langle x_0 - g_0, x_0 - g | b \rangle$$

for all $g \in K$, Thus

$$\begin{aligned} \|x - g_0, b\|^2 &\leq \inf_{g \in K} \langle x_0 - g_0, x_0 - g|b \rangle \\ &\leq \langle x_0 - g_0, x_0 - g_0|b \rangle = \|x - g_0, b\|^2 \end{aligned}$$

Therefore

$$\|x_0 - g_0, b\|^2 = \inf_{g \in K} \langle x_0 - g_0, x_0 - g|b \rangle.$$

(3) \Rightarrow (1) : If (3) holds, then by Cauchy-Schwartz inequality in 2-inner product spaces, for all $g \in K$ we have

$$\|x_0 - g_0, b\|^2 \leq \langle x_0 - g_0, x_0 - g|b \rangle \leq \|x_0 - g_0, b\| \|x_0 - g, b\|.$$

Thus $\|x_0 - g_0, b\| \leq \|x_0 - g, b\|$ for all $g \in K$. That is $g_0 = P_{K,b}(x_0)$. \square

Lemma 2.1. Let X be a 2-inner product space over the real field \mathbb{R} . Then:

(1) If S is a nonempty subset of X , then S_b° is a b -closed convex cone and S_b^{\perp} is a b -closed subspace.

(2) If C is a convex cone in X , then $(C - y)_b^\circ = C_b^\circ \cap y_b^\perp$ for each $y \in C$.

(3) If M is a subspace of X , then $M_b^\circ = M_b^\perp$.

(4) If C is a b -chebyshev convex cone in X , then $C_b^{\circ\circ} = C$.

(5) If M is a b -chebyshev subspace in X , then $M_b^{\circ\circ} = M_b^{\perp\perp} = M$.

Proof. (1) Let $x_n \in S_b^\circ$ and $x_n \xrightarrow{b} x$. Then for each $y \in S$,

$$\langle x, y|b \rangle = \lim \langle x_n, y|b \rangle \leq 0$$

implies $x \in S_b^\circ$ and S_b° is b -closed. Let $x, z \in S_b^\circ$ and $\alpha, \beta \geq 0$. Then, for each $y \in S$,

$$\langle \alpha x + \beta z, y|b \rangle = \alpha \langle x, y|b \rangle + \beta \langle z, y|b \rangle \leq 0$$

so $\alpha x + \beta z \in S_b^\circ$ and S_b° is a convex cone. Similarly we can prove S_b^\perp is a b -closed subspace.

(2) We have $x \in (C - y)_b^\circ$ if and only if $\langle x, c - y|b \rangle \leq 0$ for all $c \in C$. Taking $c = 0$ and $c = 2y$, it follows that the last statement is equivalent to $\langle x, y|b \rangle = 0$ and $\langle x, c|b \rangle \leq 0$ for all $c \in C$. That is, $x \in C_b^\circ \cap y_b^\perp$.

(3) If M is a subspace, then $-M = M$ implies

$$M_b^\circ = M_b^\circ \cap (-M)_b^\circ = M_b^\perp.$$

(4) Let C be a b -chebyshev convex cone, and $x \in C \setminus \langle b \rangle$. Then for any $y \in C_b^\circ$, $\langle x, y|b \rangle \leq 0$. Hence $x \in C_b^{\circ\circ}$. That is $C \subseteq C_b^{\circ\circ}$. Now remains to verify $C_b^{\circ\circ} \subseteq C$. If not, choose $x \in C_b^{\circ\circ} \setminus C$ and let $y_0 \in P_{C,b}(x)$. By (2) and theorem (2.1) we have

$$x - y_0 \in (C - y_0)_b^\circ = C_b^\circ \cap y_{0b}^\perp.$$

Thus

$$0 < \|x - y_0, b\|^2 = \langle x - y_0, x - y_0|b \rangle = \langle x - y_0, x|b \rangle \leq 0$$

which is absurd. Therefore $C_b^{\circ\circ} \subseteq C$.

(5) It is clear by (3),(4). \square

Now we investigate theorem (2.1) in the case of b -closed convex cone.

Theorem 2.2. Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space, C a b -closed convex cone in X with $X \neq C$. If $x_0 \in X \setminus (C + \langle b \rangle)$ and $g_0 \in C$, then the following statements are equivalent:

(1) $g_0 = P_{C,b}(x_0)$,

- (2) $x_0 - g_0 \in C_b^\circ \cap g_0^\perp_b$,
- (3) $x_0 - g_0 \in C_b^\circ$ and $\langle x_0, g_0 | b \rangle = \|g_0, b\|^2$

Proof. The equivalence of (2) and (3) is clear. The equivalence of (1) and (2) is also clear by lemma (2.2). □

Remark 2.2. If M is a linear subspace then by lemma(2.2), $M_b^\circ = M_b^\perp$ and so condition (2) of above theorem reduces to the classical condition that

$$g_0 = P_{M,b}(x_0) \iff x - g_0 \perp_b M \iff \langle x - g_0, g | b \rangle = 0 \text{ for all } g \in M ; (\text{see [13]}).$$

Remark 2.3. Let V be an affine set in the 2-inner product space X , i.e. $V = M + v$, where M is a subspace and v is any element of V . Then we have that

$$g_0 = P_{V,b}(x_0) \iff x_0 - g_0 \perp_b M \iff \langle x_0 - g_0, g - v | b \rangle = 0 \text{ for all } g \in V.$$

Moreover

$$P_{V,b}(x + e) = P_{V,b}(x) \text{ for all } x \in X, e \in M_b^\perp.$$

If $(X, \langle \cdot, \cdot | \cdot \rangle)$ is an inner product space, then the function

$$\langle x, y | z \rangle = \frac{\langle x, y \rangle \langle x, z \rangle}{\langle y, z \rangle \langle z, z \rangle} = \|z\|^2 \langle x, y \rangle - \langle x, z \rangle \langle y, z \rangle$$

for all $x, y, z \in X$ defined a 2-inner product on $X \times X \times X$.

Example 2.4. Let $X = \mathbb{R}^2$, $C = \{(y_1, y_2) \in \mathbb{R}^2; |y_2| \leq y_1\}$ and $b = (b_1, b_2) \in \mathbb{R}^2 \setminus \{(0, 0)\}$. It is simple to verify that C is a b-closed convex cone. Now if $(x_1, x_2) \in C + \langle b \rangle$, then

$$P_{C,b}(x_1, x_2) = \begin{cases} (\{ \frac{x_2 b_1 - x_1 b_2}{b_2 - b_1}, \frac{x_2 b_1 - x_1 b_2}{b_2 - b_1} \} + \langle b \rangle) \cap C & \text{if } x_2 > 0, b_1 \neq b_2 \\ ((x_1, x_2) + \langle b \rangle) \cap C & \text{if } b_1 = b_2 \\ (\{ \frac{x_2 b_1 - x_1 b_2}{b_2 + b_1}, \frac{x_1 b_2 - x_2 b_1}{b_2 + b_1} \} + \langle b \rangle) \cap C & \text{if } x_2 < 0, b_1 \neq -b_2 \\ ((x_1, x_2) + \langle b \rangle) \cap C & \text{if } b_1 = -b_2 \end{cases}$$

and if $(x_1, x_2) \in \mathbb{R}^2 \setminus (C + \langle b \rangle)$, then $P_{C,b}(x_1, x_2) = \{(0, 0)\}$.

Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space over the real field \mathbb{R} and let $F : X \times \langle b \rangle \rightarrow \mathbb{R}$ be a continues convex 2-functional on $X \times \langle b \rangle$. Denote by

$$A_b(r) := \{x \in X | F(x, b) \leq r\}$$

Assume that r is a real number that $A_b(r) \neq \emptyset$. It is clear that $A_b(r)$ is a b-closed convex subset of X .

Theorem 2.3. Let $(X, \langle \cdot, \cdot | \cdot \rangle)$ be a 2-inner product space over the real field \mathbb{R} and let $F : X \times \langle b \rangle \rightarrow \mathbb{R}$ be a convex 2-functional on $X \times \langle b \rangle$. Let $x_0 \in X \setminus (A_b(r) + \langle b \rangle)$ and $g_0 \in A_b(r)$. Then the following are equivalent.

- (1) $g_0 = P_{A_b(r),b}(x_0)$;
- (2) $F(x, b) \geq r + \frac{F(x_0,b) - r}{\|x_0 - g_0, b\|^2} \langle x - g_0, x_0 - g_0 | b \rangle$ for all $x \in A_b(r)$ where $r = F(g_0, b)$.

Proof. (1) \Rightarrow (2) Assume that $g_0 = P_{A_b(r),b}(x_0)$. Since $x_0 \in X \setminus (A_b(r) + \langle b \rangle)$ we have $F(x_0, b) > r$. Let $x \in A_b(r)$, then $F(x, b) \leq r$. Set $\alpha = F(x_0, b) - r$, $\beta = r - F(x, b)$. Then $\alpha > 0$, $\beta \geq 0$ and $0 < \alpha + \beta = F(x_0, b) - F(x, b)$. Consider the element $u = \frac{\alpha x + \beta x_0}{\alpha + \beta}$. By convexity of F , for $y = y' = b$, $\lambda = \frac{\alpha}{\alpha + \beta}$, $\mu = 1$ and $a = b = 1$ in definition of convex 2-functional we have that:

$$\begin{aligned} F(u, b) &= F\left(\frac{\alpha x + \beta x_0}{\alpha + \beta}, b\right) \leq \frac{\alpha F(x, b) + \beta F(x_0, b)}{\alpha + \beta} \\ &= \frac{(F(x_0, b) - r)F(x, b) + (r - F(x, b))F(x_0, b)}{F(x_0, b) - F(x, b)} = r \end{aligned}$$

That is $u \in A_b(r)$. As $g_0 = P_{A_b(r),b}(x_0)$, we have by theorem (2.1) that $x_0 - g_0 \in (A_b(r) - g_0)_b^\circ$, so, $\langle g - g_0, x_0 - g_0 | b \rangle \leq 0$ for all $g \in A_b(r)$. In particular, $\langle u - g_0, x_0 - g_0 | b \rangle \leq 0$. That is

$$\begin{aligned} 0 &\geq \langle u - g_0, x_0 - g_0 | b \rangle = \left\langle \frac{\alpha x + \beta x_0}{\alpha + \beta} - g_0, x_0 - g_0 | b \right\rangle \\ &= \frac{1}{\alpha + \beta} \langle \alpha x + \beta x_0 - (\alpha + \beta)g_0, x_0 - g_0 | b \rangle \\ &= \frac{\alpha}{\alpha + \beta} \langle x - g_0, x_0 - g_0 | b \rangle + \frac{\beta}{\alpha + \beta} \langle x_0 - g_0, x_0 - g_0 | b \rangle \\ &= \frac{(F(x_0, b) - r)}{F(x_0, b) - F(x, b)} \langle x - g_0, x_0 - g_0 | b \rangle + \frac{(r - F(x, b))}{F(x_0, b) - F(x, b)} \|x_0 - g_0, b\|^2 \end{aligned}$$

Thus

$$F(x, b) \geq \frac{(F(x_0, b) - r)}{\|x_0 - g_0, b\|^2} \langle x - g_0, x_0 - g_0 | b \rangle$$

for all $x \in A_b(r)$. Since above theorem is true for all $x \in A_b(r)$ so for $x = g_0$ we have that $F(g_0, b) \geq r$. But since $g_0 \in A_b(r)$ we have $F(g_0, b) \leq r$. Thus $F(g_0, b) = r$.

(2) \implies (1): Assume that (2) holds. Then for all $x \in A_b(r)$,

$$0 \geq F(x, b) - r \geq \frac{(F(x_0, b) - r)}{\|x_0 - g_0, b\|^2} \langle x - g_0, x_0 - g_0 | b \rangle$$

Since $F(x_0, b) - r > 0$, we have that

$$\langle x - g_0, x_0 - g_0 | b \rangle \leq 0$$

for all $x \in A_b(r)$. That is, $x_0 - g_0 \in (A_b(r) - g_0)^\circ$, whence, $g_0 = P_{A_b(r),b}(x_0)$. \square

Corollary 2.5. Let $f : X \times \langle b \rangle \longrightarrow \mathbb{R}$ be a continuous sublinear 2-functional on the 2-inner product space $(X, \langle \cdot, \cdot | \cdot \rangle)$. Put $K_b(f) := \{x \in X | f(x, b) \leq 0\}$. Let $x_0 \in X \setminus (K_b(f) + \langle b \rangle)$ and $g_0 \in K_b(f)$. Then the following statements are equivalent:

(1) $g_0 = P_{K_b(f),b}(x_0)$;

(2) $f(x, b) \geq \frac{f(x_0, b)}{\|x_0 - g_0, b\|^2} \langle x - g_0, x_0 - g_0 | b \rangle$ for all $x \in K_b(f)$.

Proof. It is sufficient that in above theorem taking $F = f$ and $r = 0$. \square

It is clear that $X = K_b(f) \cup K_b(-f)$ and $\ker(f) = \{x \in X | f(x, b) = 0\} = K_b(f) \cap K_b(-f)$. If in the above corollary replacing f with $-f$, then we have:

Corollary 2.6. Let $f : X \times \langle b \rangle \longrightarrow \mathbb{R}$ be a continuous sublinear 2-functional on the 2-inner product space $(X, \langle \cdot, \cdot | \cdot \rangle)$. Let $x_0 \in X \setminus (\ker(f) + \langle b \rangle)$ and $g_0 \in \ker(f)$. Then the following statements are equivalent:

(1) $g_0 = P_{\ker(f),b}(x_0)$;

(2) $f(x, b) = \frac{f(x_0, b)}{\|x_0 - g_0, b\|^2} \langle x, x_0 - g_0 | b \rangle$ for all $x \in \ker(f)$.

Remark 2.7. For another proof of above corollary see [15].

3. B-METRIC PROJECTION IN 2-INNER PRODUCT SPACES

In this section we investigate some properties of the b-metric projection onto convex cone and get some consequence, specially we show that every 2-inner product space is direct sum of any b-chebyshev subspace and its b-orthogonal complement.

Proposition 3.1. *Let K be a convex b -chebyshev set and $K \cap \langle b \rangle = \emptyset$. Then*

(1) $P_{K,b}$ is idempotent i.e.

$$P_{K,b}(P_{K,b}(x)) = P_{K,b}(x)$$

for every $x \in X$.

(2) $P_{K,b}$ is firmly nonexpansive i.e.

$$\langle x - y, P_{K,b}(x) - P_{K,b}(y) | b \rangle \geq \|P_{K,b}(x) - P_{K,b}(y), b\|^2$$

for all $x, y \in X$.

(3) $P_{K,b}$ is monotone i.e.

$$\langle x - y, P_{K,b}(x) - P_{K,b}(y) | b \rangle \geq 0$$

for all $x, y \in X$.

(4) $P_{K,b}$ is strictly nonexpansive i.e.

$$\|x - y, b\|^2 > \|P_{K,b}(x) - P_{K,b}(y), b\|^2 + \|x - P_{K,b}(x) - (y - P_{K,b}(y)), b\|^2$$

for all $x, y \in X$.

(5) $P_{K,b}$ is nonexpansive i.e.

$$\|P_{K,b}(x) - P_{K,b}(y), b\| \leq \|x - y, b\|$$

for all $x, y \in X$.

(6) $P_{K,b}$ is uniformly continuous.

Proof. (1) It is clear.

(2) We have:

$$\begin{aligned} \langle x - y, P_{K,b}(x) - P_{K,b}(y) | b \rangle &= \langle x - P_{K,b}(x), P_{K,b}(x) - P_{K,b}(y) | b \rangle \\ &\quad + \langle P_{K,b}(x) - P_{K,b}(y), P_{K,b}(x) - P_{K,b}(y) | b \rangle \\ &\quad + \langle P_{K,b}(y) - y, P_{K,b}(x) - P_{K,b}(y) | b \rangle \end{aligned}$$

The first and third terms on the right are nonnegative by theorem (2.1), and the second term is $\|P_{K,b}(x) - P_{K,b}(y), b\|^2$. This verifies (2).

(3) It is immediate consequence of (2).

(4) Using (2) we obtain for each $x, y \in X$ that:

$$\begin{aligned} \|x - y, b\|^2 &= \|(x - P_{K,b}(x)) + (P_{K,b}(x) - P_{K,b}(y)) + (P_{K,b}(y) - y), b\|^2 \\ &= \|P_{K,b}(x) - P_{K,b}(y), b\|^2 + \|(x - P_{K,b}(x)) - (y - P_{K,b}(y)), b\|^2 \\ &\quad + 2\langle P_{K,b}(x) - P_{K,b}(y), x - P_{K,b}(x) - (y - P_{K,b}(y)) | b \rangle \\ &= \|P_{K,b}(x) - P_{K,b}(y), b\|^2 + \|(x - P_{K,b}(x)) - (y - P_{K,b}(y)), b\|^2 \\ &\quad + 2\langle P_{K,b}(x) - P_{K,b}(y), x - y | b \rangle - 2\|P_{K,b}(x) - P_{K,b}(y), b\|^2 \\ &\geq \|P_{K,b}(x) - P_{K,b}(y), b\|^2 + \|x - P_{K,b}(x) - (y - P_{K,b}(y)), b\|^2 \end{aligned}$$

This proves (4).

(5) It follows immediately from (4).

(6) It follows immediately from (5). □

Theorem 3.1. *Let C be a b -chebyshev convex cone in the 2-inner product space X and $C \cap \langle b \rangle = \emptyset$. Then C_b° is a b -chebyshev convex cone and*

(1) For each $x \in X$,

$$x = P_{C,b}(x) + P_{C_b^\circ,b}(x) \text{ and } P_{C,b}(x) \perp_b P_{C_b^\circ,b}(x).$$

Moreover, this representation is unique in the sense that if $x = y + z$ for some $y \in C$ and $z \in C_b^\circ$ with $y \perp_b z$, then $y = P_{C,b}(x)$ and $z = P_{C_b^\circ,b}(x)$.

- (2) $\|x, b\|^2 = \|P_{C,b}(x), b\|^2 + \|P_{C_b^\circ,b}(x), b\|^2$ for all $x \in X \setminus \langle b \rangle$.
 (3) $C_b^\circ = \{x \in X | P_{C,b}(x) = 0\}$ and $C = \{x \in X | P_{C_b^\circ,b}(x) = 0\} = \{x \in X | P_{C,b}(x) = x\}$.
 (4) $\|P_{C,b}(x), b\| \leq \|x, b\|$ for all $x \in X$; moreover, $\|P_{C,b}(x), b\| = \|x, b\|$ if and only if $x \in C$.
 (5) $C_b^{\circ\circ} = C$.
 (6) $P_{C,b}$ is positively homogeneous. i.e.

$$P_{C,b}(\lambda x) = \lambda P_{C,b}(x) \text{ for all } x \in X, \lambda \geq 0.$$

Proof. (1) Let $x \in X$ and $c_0 = x - P_{C,b}(x)$. By theorem (2.3) $c_0 \in C_b^\circ$ and $c_0 \perp_b (x - c_0)$. For every $y \in C_b^\circ$,

$$\langle x - c_0, y | b \rangle = \langle P_{C,b}(x), y | b \rangle \leq 0.$$

Hence $x - c_0 \in (C_b^\circ)^\circ$. By theorem (2.3), we get that $c_0 = P_{C_b^\circ,b}(x)$. This proves that C_b° is b -chebyshev convex cone, $x = P_{C,b}(x) + P_{C_b^\circ,b}(x)$ and $P_{C,b}(x) \perp_b P_{C_b^\circ,b}(x)$. Now we verify the uniqueness of this representation. Let $x = y + z$, where $y \in C, z \in C_b^\circ$ and $y \perp_b z$. For each $c \in C$,

$$\langle x - y, c | b \rangle = \langle z, c | b \rangle \leq 0$$

and

$$\langle x - y, y | b \rangle = \langle z, y | b \rangle = 0.$$

By theorem (2.3) $y = P_{C,b}(x)$. Similarly $z = P_{C_b^\circ,b}(x)$.

(2) It is clear by (1) and Pythagorean theorem in 2-inner product spaces.

(3) By using (1) we have that:

$$x \in C_b^\circ \iff x = P_{C_b^\circ,b}(x) \iff P_{C,b}(x) = 0,$$

and

$$x \in C \iff x = P_{C,b}(x) \iff P_{C_b^\circ,b}(x) = 0.$$

(4) From (2), it is clear that $\|P_{C,b}(x), b\| \leq \|x, b\|$ for all $x \in X$. Also from (2), $\|P_{C,b}(x), b\| = \|x, b\|$ if and only if $P_{C_b^\circ,b}(x) = 0$, which from (3) is equivalent to $x \in C$.

(5) By (3) we have:

$$C_b^{\circ\circ} = (C_b^\circ)^\circ = \{x \in X | P_{C_b^\circ,b}(x) = 0\} = C.$$

(6) Let $x \in X$ and $\lambda \geq 0$. Then $x = P_{C,b}(x) + P_{C_b^\circ,b}(x)$ and $\lambda x = \lambda P_{C,b}(x) + \lambda P_{C_b^\circ,b}(x)$. Since both C and C_b° are convex cones, $\lambda P_{C,b}(x) \in C$ and $\lambda P_{C_b^\circ,b}(x) \in C_b^\circ$. By (1) and the uniqueness of representation for λx , we see that $P_{C,b}(\lambda x) = \lambda P_{C,b}(x)$. \square

Corollary 3.2. Let M be a b -chebyshev subspace of the 2-inner product space X . Then M_b^\perp is a b -chebyshev subspace and:

(1) $x = P_{M,b}(x) + P_{M_b^\perp,b}(x)$, for each $x \in X$. Moreover, this representation is unique in the sense that if $x = y + z$ where $y \in M$ and $z \in M_b^\perp$, then $y = P_{M,b}(x)$ and $z = P_{M_b^\perp,b}(x)$.

(2) $\|x, b\|^2 = \|P_{M,b}(x), b\|^2 + \|P_{M_b^\perp,b}(x), b\|^2$ for all $x \in X \setminus \langle b \rangle$.

(3) $M_b^\perp = \{x \in X | P_{M,b}(x) = 0\}$ and $M = \{x \in X | P_{M_b^\perp,b}(x) = 0\} = \{x \in X | P_{M,b}(x) = x\}$.

(4) $\|P_{M,b}(x), b\| \leq \|x, b\|$ for all $x \in X$; moreover, $\|P_{M,b}(x), b\| = \|x, b\|$ if and only if $x \in M$.

(5) $M_b^{\perp\perp} = M$.

Theorem 3.2. Let M be a b -chebyshev subspace of the 2-inner product space X and $M \cap \langle b \rangle = \emptyset$. Then:

(1) $P_{M,b}$ is a bounded linear 2-functional and $\|P_{M,b}\| = 1$ (in the case $M = \{0\}$, we have $\|P_{M,b}\| = 0$).

(2) $P_{M,b}$ is self-adjoint i.e.

$$\langle P_{M,b}(x), y|b \rangle = \langle x, P_{M,b}(y)|b \rangle \text{ for all } x, y \in X.$$

(3) For every $x \in X$,

$$\langle P_{M,b}(x), x|b \rangle = \|P_{M,b}(x), b\|^2.$$

(4) $P_{M,b}$ is nonnegative i.e.

$$\langle P_{M,b}(x), x|b \rangle \geq 0 \text{ for every } x \in X.$$

Proof. (1) Let $x, y \in X$ and $\alpha, \beta \in \mathbb{R}$. By remark (2.4), $x - P_{M,b}(x)$ and $y - P_{M,b}(y)$ are in M_b^\perp . since M_b^\perp is subspace,

$$\alpha x + \beta y - (\alpha P_{M,b}(x) + \beta P_{M,b}(y)) = \alpha(x - P_{M,b}(x)) + \beta(y - P_{M,b}(y)) \in M_b^\perp.$$

Since $\alpha P_{M,b}(x) + \beta P_{M,b}(y) \in M$, remark (2.4) implies that $\alpha P_{M,b}(x) + \beta P_{M,b}(y) = P_{M,b}(\alpha x + \beta y)$. Thus $P_{M,b}$ is linear. From corollary (3.3 [4]) we get $\|P_{M,b}(x)\| \leq \|x, b\|$ for all $x \in X$. So $\|P_{M,b}$ is bounded and $\|P_{M,b}\| \leq 1$. Since $P_{M,b}y = y$ for all $y \in M$ and $\|y, b\| = \|P_{M,b}(y)\| \leq \|P_{M,b}\| \|y, b\|$, implies that $\|P_{M,b}\| \geq 1$ therefore $\|P_{M,b}\| = 1$.

(2) By remark (2.4), for each $x, y \in X$ we have $\langle P_{M,b}(x), y - P_{M,b}(y)|b \rangle = 0$, and hence

$$\langle P_{M,b}(x), y|b \rangle = \langle P_{M,b}(x), P_{M,b}(y)|b \rangle. (*)$$

By replacing x and y in above relation we obtain

$$\begin{aligned} \langle x, P_{M,b}(y)|b \rangle &= \langle P_{M,b}(y), x|b \rangle \\ &= \langle P_{M,b}(y), P_{M,b}(x)|b \rangle \\ &= \langle P_{M,b}(x), P_{M,b}(y)|b \rangle \\ &= \langle P_{M,b}(x), y|b \rangle. \end{aligned}$$

(3) Tacking $y = x$ in (*).

(4) It is clear from (3). □

REFERENCES

- [1] Y. J. Cho, P. C. S. Lin, S. S. Kim and A. Misiak, Theory of 2-inner product spaces, *New York: Nova Science Publishes, Inc.*, 2001.
- [2] S. S. Dragomir, Best approximation in pre-Hilbertian spaces and applications to continuous sub-linear functionals, *Anal. Uni. Timisoara, Ser St. Mat.*, 22, 137-140, 1991.
- [3] R. E. Ehret, Linear 2-normed spaces, *Doctoral Dissertation, St. Louis University*, 1969.
- [4] S. Elumalai, Best approximation sets in 2-normed spaces, *Comm. Korean. Math. Soc.*, 12, 619-629, 1997.
- [5] S. Gähler, Lineare 2-normierte Räume, *Math Nachr.*, 28: 1-43, 1964.
- [6] S. S. Kim and M. Crasmareanu, Best approximations and orthogonalites in 2k-inner product spaces, *Bull. Korean. Math. Soc.*, 43(2): 377-387, 2006.
- [7] S. N. Lal, Mohan Das, 2-functionals and some extention theorems in linear spaces, *Indian. J. PureAppl. Math.*, 13(8): 912-919, 1982.
- [8] Z. Lewandowska, Generalized 2-normed spaces, *Supskie Space Matemayczno Fizyczne* 1, 33-40, 2001.
- [9] H. Mazaheri and R. Kazemi, Some results on 2-inner product spaces, *Novi Sad J. Math.*, 37(2): 35-40, 2007.
- [10] A. Misiak, n-inner product spaces, *Math. Nachr.*, 140, 299-319, 1989.
- [11] T. D. Narang, Best approximation on convex sets in metric linear spaces, *Math. Nachr.*, 78, 125-130, 1997.

- [12] R. Ravi, Approximation in linear 2-normed spaces and normed linear spaces., *Doctoral Thesis, Madras Univ.*, 1994.
- [13] Sh. Rezapour, Proximinal subspaces of 2-normed spaces, *Anal. Theory. Appl.*, 22(2), 114-119, 2006.
- [14] I. Singer, Best Approximation in normed linear space by element of linear subspaces, *Springer, Berlin*, 1970.
- [15] T. Sistani and M. Abrishami Moghaddam, Some results on best approximation in convex subset of 2-normed spaces, *Int. J. Math. Analysis*, 21(3): 1043-1049, 2009.