

SOME COMMON FIXED POINT RESULTS FOR GENERALIZED WEAK CONTRACTIVE MAPPINGS IN PARTIALLY ORDERED METRIC SPACES

SUMIT CHANDOK*

Department of Mathematics, Khalsa College of Engineering Technology (Punjab Technical University),
Ranjit Avenue, Amritsar-143001, India

ABSTRACT. Some common fixed point results satisfying a generalized weak contractive condition in the framework of partially ordered metric spaces are obtained. The proved results generalize and extend some known results in the literature.

KEYWORDS : Common fixed point; f -Weakly contractive mappings; (μ, ψ) -Generalized f -weakly contractive mappings; Weakly compatible mappings

AMS Subject Classification: 41A50 47H10 54H25

1. INTRODUCTION AND PRELIMINARIES

Fixed point theory is an old and rich branch of analysis and has a large number of applications. Fixed point problems involving different contractive type inequalities have been studied by many authors (see [1]-[20] and references cited therein). The main aim of this work is to prove some common fixed point theorems for (μ, ψ) -generalized f -weakly contractive mappings in partially ordered metric spaces.

The Banach contraction mapping is one of the pivotal results of analysis. It is very popular tool for solving existence problems in many different fields of mathematics. There are a lot of generalizations of the Banach contraction principle in the literature. Ran and Reurings [18] extended the Banach contraction principle in partially ordered sets with some applications to linear and nonlinear matrix equations. While Nieto and Rodríguez-López [17] extended the result of Ran and Reurings and applied their main theorems to obtain a unique solution for a first order ordinary differential equation with periodic boundary conditions. Bhaskar and Lakshmikantham [2] introduced the concept of mixed monotone mappings and obtained some coupled fixed point results. Also, they applied their results on a first order differential equation with periodic boundary conditions.

* Corresponding author.

Email address : chansok.s@gmail.com, chandhok.sumit@gmail.com(Sumit Chandok).

Article history : Received 29 July 2012. Accepted 1 October 2012.

Alber and Guerre-Delabriere [1] introduced the concept of weakly contractive mappings and proved the existence of fixed points for single-valued weakly contractive mappings in Hilbert spaces. Thereafter, in 2001, Rhoades [20] proved the fixed point theorem which is one of the generalizations of Banach's Contraction Mapping Principle, because the weakly contractions contains contractions as a special case and he also showed that some results of [1] are true for any Banach space. In fact, weakly contractive mappings are closely related to the mappings of Boyd and Wong [3] and of Reich types [19]. Fixed point problems involving weak contractions and mappings satisfying weak contractive type inequalities have been studied by many authors (see [1], [7]-[15], [20] and references cited therein).

First, we recall some basic definitions and related results.

A map $T : X \longrightarrow X$ is called a *weakly contractive mapping* (see [1], [13], [20]) if for each $x, y \in X$,

$$d(Tx, Ty) \leq d(x, y) - \psi(d(x, y)) \quad (1.1)$$

where $\psi : [0, \infty) \longrightarrow [0, \infty)$ is continuous and nondecreasing, $\psi(x) = 0$ if and only if $x = 0$ and $\lim \psi(x) = \infty$.

If we take $\psi(x) = kx$, $0 < k < 1$, then a weakly contractive mapping is called a contraction.

A map $T : X \longrightarrow X$ is called a *f-weakly contractive mapping* (see [14]) if for each $x, y \in X$,

$$d(Tx, Ty) \leq d(fx, fy) - \psi(d(fx, fy)) \quad (1.2)$$

where $f : X \longrightarrow X$ is a self-mapping, $\psi : [0, \infty) \longrightarrow [0, \infty)$ is continuous and nondecreasing, $\psi(x) = 0$ if and only if $x = 0$ and $\lim \psi(x) = \infty$.

If we take $\psi(x) = (1 - k)x$, $0 < k < 1$, then a *f-weakly contractive mapping* is called a *f-contraction*. Further, if $f =$ identity mapping and $\psi(x) = (1 - k)x$, $0 < k < 1$, then a *f-weakly contractive mapping* is called a contraction.

A map $T : X \longrightarrow X$ is called a *generalized f-weakly contractive mapping* (see [7]) if for each $x, y \in X$,

$$d(Tx, Ty) \leq \frac{1}{2}[d(fx, Ty) + d(fy, Tx)] - \psi(d(fx, Ty), d(fy, Tx)) \quad (1.3)$$

where $f : X \longrightarrow X$ is a self-mapping, $\psi : [0, \infty)^2 \longrightarrow [0, \infty)$ is a continuous mapping such that $\psi(x, y) = 0$ if and only if $x = y = 0$.

If $f =$ identity mapping, then a generalized *f-weakly contractive mapping* is a generalized weakly contractive mapping (see [13]).

Khan et al. [16] initiated the use of a control function that alters distance between two points in a metric space, which they called an altering distance function.

A function $\mu : [0, \infty) \longrightarrow [0, \infty)$ is called an *altering distance function* if the following properties are satisfied:

- (i) μ is monotone increasing and continuous;
- (ii) $\mu(t) = 0$ if and only if $t = 0$.

A map $T : X \longrightarrow X$ is called a (μ, ψ) -*generalized f-weakly contractive mapping* (see [8]) if for each $x, y \in X$,

$$\mu(d(Tx, Ty)) \leq \mu\left(\frac{1}{2}[d(fx, Ty) + d(fy, Tx)]\right) - \psi(d(fx, Ty), d(fy, Tx)) \quad (1.4)$$

where $f : X \longrightarrow X$ is a self-mapping, $\mu : [0, \infty) \longrightarrow [0, \infty)$ is an altering distance function and $\psi : [0, \infty)^2 \longrightarrow [0, \infty)$ is a lower semi-continuous mapping such that $\psi(x, y) = 0$ if and only if $x = y = 0$.

If $f =$ identity mapping, then a (μ, ψ) -*generalized f-weakly contractive mapping* is a (μ, ψ) -*generalized weakly contractive mapping*.

Let M be a nonempty subset of a metric space (X, d) , a point $x \in M$ is a *common fixed (coincidence) point* of f and T if $x = fx = Tx$ ($fx = Tx$). The set of fixed points (respectively, coincidence points) of f and T is denoted by $F(f, T)$ (respectively, $C(f, T)$). The mappings $T, f : M \rightarrow M$ are called *commuting* if $Tfx = fTx$ for all $x \in M$; *compatible* if $\lim d(Tfx_n, fTx_n) = 0$ whenever $\{x_n\}$ is a sequence such that $\lim Tx_n = \lim fx_n = t$ for some t in M ; *weakly compatible* if they commute at their coincidence points, i.e., if $fTx = Tfx$ whenever $fx = Tx$.

Suppose (X, \leq) is a partially ordered set and $T, f : X \rightarrow X$. A mapping T is said to be *monotone f -nondecreasing* if for all $x, y \in X$,

$$fx \leq fy \text{ implies } Tx \leq Ty. \quad (1.5)$$

If f =identity mapping, then T is a *monotone nondecreasing*.

A subset W of a partially ordered set X is said to be *well ordered* if every two elements of W are comparable.

2. MAIN RESULTS

Theorem 2.1. *Let (X, \leq) be a partially ordered set and suppose that there exists a metric d on X such that (X, d) is a complete metric space. Suppose that T and f are self mappings on X , $T(X) \subseteq f(X)$, T is a monotone f -nondecreasing mapping and*

$$\mu(d(Tx, Ty)) \leq \mu\left(\frac{1}{2}[d(fx, Ty) + d(fy, Tx)]\right) - \psi(d(fx, Ty), d(fy, Tx)) \quad (2.1)$$

for all $x, y \in X$ for which $fx \geq fy$ where $\mu : [0, \infty) \rightarrow [0, \infty)$ is an altering distance function and $\psi : [0, \infty)^2 \rightarrow [0, \infty)$ is a lower semi-continuous mapping such that $\psi(x, y) = 0$ if and only if $x = y = 0$.

If $\{f(x_n)\} \subset X$ is a nondecreasing sequence with $f(x_n) \rightarrow f(z)$ in $f(X)$, then $f(x_n) \leq f(z)$, and $f(z) \leq f(f(z))$ for every n .

Also suppose that $f(X)$ is closed. If there exists an $x_0 \in X$ with $f(x_0) \leq T(x_0)$, then T and f have a coincidence point.

Further, if T and f are weakly compatible, then T and f have a common fixed point. Moreover, the set of common fixed points of T and f is well ordered if and only if T and f have one and only one common fixed point.

Proof. Let $x_0 \in X$ such that $f(x_0) \leq T(x_0)$. Since $T(X) \subseteq f(X)$, we can choose $x_1 \in X$ so that $fx_1 = Tx_0$. Since $Tx_1 \in f(X)$, there exists $x_2 \in X$ such that $fx_2 = Tx_1$. By induction, we construct a sequence $\{x_n\}$ in X such that $fx_{n+1} = Tx_n$, for every $n \geq 0$.

Since $f(x_0) \leq T(x_0)$, $T(x_0) = f(x_1)$, $f(x_0) \leq f(x_1)$, T is monotone f -nondecreasing mapping, $T(x_0) \leq T(x_1)$. Similarly $f(x_1) \leq f(x_2)$, $T(x_1) \leq T(x_2)$, $f(x_2) \leq f(x_3)$. Continuing, we obtain

$$T(x_0) \leq T(x_1) \leq T(x_2) \leq \dots \leq T(x_n) \leq T(x_{n+1}) \leq \dots$$

We suppose that $d(T(x_n), T(x_{n+1})) > 0$ for all n . If not then $T(x_{n+1}) = T(x_n)$ for some n , $T(x_{n+1}) = f(x_{n+1})$, i.e. T and f have a coincidence point x_{n+1} , and so we have the result.

Consider

$$\begin{aligned} \mu(d(Tx_{n+1}, Tx_n)) &\leq \mu\left(\frac{1}{2}[d(fx_{n+1}, Tx_n) + d(fx_n, Tx_{n+1})]\right) - \psi(d(fx_{n+1}, Tx_n), d(fx_n, Tx_{n+1})) \\ &= \mu\left(\frac{1}{2}d(Tx_{n-1}, Tx_{n+1})\right) - \psi(0, d(Tx_{n-1}, Tx_{n+1})) \\ &\leq \mu\left(\frac{1}{2}d(Tx_{n-1}, Tx_{n+1})\right) \end{aligned} \quad (*)$$

$$\leq \mu\left(\frac{1}{2}[d(Tx_{n-1}, Tx_n) + d(Tx_n, Tx_{n+1})]\right)$$

Since μ is a non-decreasing function, for all $n = 1, 2, \dots$, we have $d(Tx_{n+1}, Tx_n) \leq d(Tx_n, Tx_{n-1})$. Thus $\{d(Tx_{n+1}, Tx_n)\}$ is a monotone decreasing sequence of non-negative real numbers and hence is convergent. Hence there exists $r \geq 0$ such that $d(Tx_{n+1}, Tx_n) \longrightarrow r$.

From inequality (*), we have

$$\begin{aligned} d(Tx_{n+1}, Tx_n) &\leq \frac{1}{2}d(Tx_{n-1}, Tx_{n+1}) \\ &\leq \frac{1}{2}[d(Tx_{n-1}, Tx_n) + d(Tx_n, Tx_{n+1})] \end{aligned}$$

letting $n \longrightarrow \infty$, we have

$$r \leq \lim \frac{1}{2}d(Tx_{n-1}, Tx_{n+1}) \leq \frac{1}{2}r + \frac{1}{2}r,$$

i.e. $\lim d(Tx_{n-1}, Tx_{n+1}) = 2r$. Using the continuity of μ and lower semi-continuity of ψ , and inequality (*), we have $\mu(r) \leq \mu(r) - \psi(0, 2r)$, and consequently, $\psi(0, 2r) \leq 0$. Thus $r = 0$. Hence

$$d(Tx_{n+1}, Tx_n) \longrightarrow 0.$$

Now, we show that $\{Tx_n\}$ is a Cauchy sequence. If otherwise, then there exists $\epsilon > 0$ for which we can find subsequences $\{Tx_{m(k)}\}$ and $\{Tx_{n(k)}\}$ of $\{Tx_n\}$ with $n(k) > m(k) > k$ such that for every k , $d(Tx_{m(k)}, Tx_{n(k)}) \geq \epsilon$, $d(Tx_{m(k)}, Tx_{n(k)-1}) < \epsilon$. So, we have

$$\begin{aligned} \epsilon &\leq d(Tx_{m(k)}, Tx_{n(k)}) \\ &\leq d(Tx_{m(k)}, Tx_{n(k)-1}) + d(Tx_{n(k)-1}, Tx_{n(k)}) \\ &< \epsilon + d(Tx_{n(k)-1}, Tx_{n(k)}). \end{aligned}$$

Letting $k \longrightarrow \infty$ and using $d(Tx_{n+1}, Tx_n) \longrightarrow 0$, we have

$$\lim d(Tx_{m(k)}, Tx_{n(k)}) = \epsilon = \lim d(Tx_{m(k)}, Tx_{n(k)-1}). \quad (2.2)$$

Again,

$$d(Tx_{m(k)}, Tx_{n(k)-1}) \leq d(Tx_{m(k)}, Tx_{m(k)-1}) + d(Tx_{m(k)-1}, Tx_{n(k)}) + d(Tx_{n(k)}, Tx_{n(k)-1}),$$

and

$$d(Tx_{m(k)-1}, Tx_{n(k)}) \leq d(Tx_{m(k)-1}, Tx_{m(k)}) + d(Tx_{m(k)}, Tx_{n(k)}).$$

Letting $k \longrightarrow \infty$ in the above two inequalities and using (2.2) we get,

$$\lim d(Tx_{m(k)-1}, Tx_{n(k)}) = \epsilon.$$

Also, we have

$$\begin{aligned} \mu(\epsilon) &\leq \mu(d(Tx_{m(k)}, Tx_{n(k)})) \\ &\leq \mu\left(\frac{1}{2}[d(Tx_{m(k)}, Tx_{n(k)}) + d(Tx_{n(k)}, Tx_{m(k)})]\right) - \\ &\quad \psi(d(Tx_{m(k)}, Tx_{n(k)}), d(Tx_{n(k)}, Tx_{m(k)})) \\ &= \mu\left(\frac{1}{2}[d(Tx_{m(k)-1}, Tx_{n(k)}) + d(Tx_{n(k)-1}, Tx_{m(k)})]\right) - \\ &\quad \psi(d(Tx_{m(k)-1}, Tx_{n(k)}), d(Tx_{n(k)-1}, Tx_{m(k)})). \end{aligned}$$

Taking $k \longrightarrow \infty$, and using the continuity of μ and lower semi-continuity of ψ , we have $\mu(\epsilon) \leq \mu(\frac{1}{2}[\epsilon + \epsilon]) - \psi(\epsilon, \epsilon)$ and consequently $\psi(\epsilon, \epsilon) \leq 0$, which is contradiction since $\epsilon > 0$. Thus $\{Tx_n\}$ is a Cauchy sequence. As $f(X)$ is closed

and $fx_n = Tx_{n-1}$, $\{fx_n\}$ is also a Cauchy sequence, there is some $z \in X$ such that $\lim fx_{n+1} = \lim Tx_n = fz$. Since $\{f(x_n)\}$ is a nondecreasing sequence and $\lim fx_{n+1} = fz$, $f(x_n) \leq f(z)$, and $f(z) \leq f(f(z))$ for every n . Consider

$$\begin{aligned} \mu(d(Tz, fx_{n+1})) &= \mu(d(Tz, Tx_n)) \\ &\leq \mu\left(\frac{1}{2}[d(fz, Tx_n) + d(fx_n, Tz)]\right) - \psi(d(fz, Tx_n), d(fx_n, Tz)), \end{aligned}$$

letting $n \rightarrow \infty$, we have

$$\mu(d(Tz, fz)) \leq \mu\left(\frac{1}{2}d(fz, Tz)\right) - \psi(0, d(fz, Tz))$$

This implies that $d(Tz, fz) = 0$, i.e. $Tz = fz$ and z is a coincidence point of T and f .

Now suppose that T and f are weakly compatible. Let $w = T(z) = f(z)$. Then $T(w) = T(f(z)) = f(T(z)) = f(w)$ and $f(z) \leq f(f(z)) = f(w)$. Consider

$$\begin{aligned} \mu(d(T(z), T(w))) &\leq \mu\left(\frac{1}{2}[d(fz, Tw) + d(fw, Tz)]\right) - \psi(d(fz, Tw), d(fw, Tz)) \\ &= \mu\left(\frac{1}{2}[d(Tz, Tw) + d(Tw, Tz)]\right) - \psi(d(Tz, Tw), d(Tw, Tz)) \\ &= \mu(d(Tw, Tz)) - \psi(d(Tz, Tw), d(Tw, Tz)). \end{aligned}$$

This implies that $d(Tz, Tw) = 0$, by the property of ψ . Therefore, $T(w) = f(w) = w$.

Now suppose that the set of common fixed points of T and f is well ordered. We claim that common fixed points of T and f is unique. Assume on contrary that, $Tu = fu = u$ and $Tv = fv = v$ but $u \neq v$. Consider

$$\begin{aligned} \mu(d(u, v)) &= \mu(d(Tu, Tv)) \\ &\leq \mu\left(\frac{1}{2}[d(fu, Tv) + d(fv, Tu)]\right) - \psi(d(fu, Tv), d(fv, Tu)) \\ &= \mu\left(\frac{1}{2}[d(u, v) + d(v, u)]\right) - \psi(d(u, v), d(v, u)) \\ &= \mu(d(u, v)) - \psi(d(u, v), d(u, v)). \end{aligned}$$

This implies that $d(u, v) = 0$, by the property of ψ . Hence $u = v$. Conversely, if T and f have only one common fixed point then the set of common fixed point of f and T being singleton is well ordered. \square

If $f = \text{identity mapping}$, then we have the following result.

Corollary 2.2. *Let (X, \leq) be a partially ordered set and suppose that there exists a metric d on X such that (X, d) is a complete metric space. Suppose that T is a self mapping on X , T is a monotone nondecreasing mapping and*

$$\mu(d(Tx, Ty)) \leq \mu\left(\frac{1}{2}[d(x, Ty) + d(y, Tx)]\right) - \psi(d(x, Ty), d(y, Tx)) \quad (2.3)$$

for all $x, y \in X$ for which $x \geq y$ where $\mu : [0, \infty) \rightarrow [0, \infty)$ is an altering distance function and $\psi : [0, \infty)^2 \rightarrow [0, \infty)$ is a lower semi-continuous mapping such that $\psi(x, y) = 0$ if and only if $x = y = 0$.

Also suppose that either

(i) $\{x_n\} \subset X$ is a nondecreasing sequence with $x_n \rightarrow z$ in X , then $x_n \leq z$, for every n ; or

(ii) T is continuous.

If there exists an $x_0 \in X$ with $x_0 \leq T(x_0)$, then T has a fixed point.

Moreover, for arbitrary two points $x, y \in X$, there exists $w \in X$ such that w is comparable with both x and y . Then the fixed point of T is unique.

Proof. If (i) holds, then taking $f = \text{identity mapping}$ in Theorem 2.1 we get the result.

If (ii) holds then proceeding as in Theorem 2.1 with $f = \text{identity mapping}$, we can prove that $\{Tx_n\}$ is a cauchy sequence, $z = \lim x_{n+1} = \lim T(x_n) = T(\lim x_n) = T(z)$ and hence T has a fixed point.

Let u and v be two fixed points of T such that $u \neq v$. Now, consider the following two cases:

(a) If u and v are comparable. Consider

$$\begin{aligned} \mu(d(u, v)) &= \mu(d(Tu, Tv)) \\ &\leq \mu\left(\frac{1}{2}[d(u, Tv) + d(v, Tu)]\right) - \psi(d(u, Tv), d(v, Tu)) \\ &\leq \mu\left(\frac{1}{2}[d(u, v) + d(v, u)]\right) - \psi(d(u, v), d(v, u)) \\ &= \mu(d(u, v)) - \psi(d(u, v), d(u, v)). \end{aligned}$$

This implies that $d(u, v) = 0$, by the property of ψ . Hence $u = v$.

(b) If u and v are not comparable. Choose an element $w \in X$ comparable with both of them. Then also $u = T^n u$ is comparable with $T^n w$ for each n . Consider

$$\begin{aligned} \mu(d(u, T^n w)) &= \mu(d(T^n u, T^n w)) \\ &= \mu(d(TT^{n-1}u, TT^{n-1}w)) \\ &\leq \mu\left(\frac{1}{2}[d(T^{n-1}u, T^n w) + d(T^{n-1}w, T^n u)]\right) - \psi(d(T^{n-1}u, T^n w), d(T^{n-1}w, T^n u)) \\ &= \mu\left(\frac{1}{2}[d(u, T^n w) + d(T^{n-1}w, u)]\right) - \psi(d(u, T^n w), d(T^{n-1}w, u)) \quad (**) \\ &\leq \mu\left(\frac{1}{2}[d(u, T^n w) + d(T^{n-1}w, u)]\right) \end{aligned}$$

and hence we get $d(u, T^n w) \leq d(u, T^{n-1}w)$. This proves that the nonnegative decreasing sequence $\{d(u, T^n w)\}$ is convergent. If $\lim_{n \rightarrow \infty} \{d(u, T^n w)\} = r$, then, letting $n \rightarrow \infty$ in $(**)$ and from the continuity of μ and lower semi-continuity of ψ we obtain $\mu(r) \leq \mu(r) - \psi(r, r) \leq \mu(r)$. This gives $\psi(r, r) = 0$ and by our assumption about ψ , $r = 0$. Consequently, $\lim_{n \rightarrow \infty} d(u, T^n w) = 0$. Analogously, it can be proved that $\lim_{n \rightarrow \infty} d(v, T^n w) = 0$. Since the limit is unique, we have $u = v$. □

If $\mu(t) = t$, then we have the following result.

Corollary 2.3. *Let (X, \leq) be a partially ordered set and suppose that there exists a metric d on X such that (X, d) is a complete metric space. Suppose that T is a self mapping on X , T is a monotone nondecreasing mapping and*

$$d(Tx, Ty) \leq \frac{1}{2}[d(x, Ty) + d(y, Tx)] - \psi(d(x, Ty), d(y, Tx)) \quad (2.4)$$

for all $x, y \in X$ for which $x \geq y$ where $\mu : [0, \infty) \rightarrow [0, \infty)$ is an altering distance function and $\psi : [0, \infty)^2 \rightarrow [0, \infty)$ is a lower semi-continuous mapping such that $\psi(x, y) = 0$ if and only if $x = y = 0$.

Also suppose that either

(i) $\{x_n\} \subset X$ is a nondecreasing sequence with $x_n \rightarrow z$ in X , then $x_n \leq z$, for every n ; or

(ii) T is continuous.

If there exists an $x_0 \in X$ with $x_0 \leq T(x_0)$, then T has a fixed point.

Moreover, for arbitrary two points $x, y \in X$, there exists $w \in X$ such that w is comparable with both x and y . Then the fixed point of T is unique.

If $\psi(x, y) = (\frac{1}{2} - k)(x + y)$, $0 < k < \frac{1}{2}$, we have the following result.

Corollary 2.4. [7] Let (X, \leq) be a partially ordered set and suppose that there exists a metric d on X such that (X, d) is a complete metric space. Suppose that T is a nondecreasing self-mapping of X and T satisfies

$$d(Tx, Ty) \leq k[d(x, Ty) + d(y, Tx)], \text{ for } x \geq y, \quad (2.5)$$

where $0 < k < \frac{1}{2}$, for all $x, y \in X$. Also suppose either

(i) if $\{x_n\} \subset X$ is a nondecreasing sequence with $x_n \longrightarrow z$ in X , then $x_n \leq z$ for every n .

or

(ii) T is continuous.

If there exists an $x_0 \in X$ with $x_0 \leq T(x_0)$, then T has a fixed point.

Acknowledgement. The author is thankful to the learned referee for valuable suggestions leading to an improvement of the paper.

REFERENCES

- [1] Ya. I. Alber and S. Guerre-Delabriere, Principles of weakly contractive maps in Hilbert spaces, New results in Operator Theory, Advances Appl., Vol. 8 (I. Gohberg and Yu. Lyubich, eds.), Birkhauser, Basel, 1997, 7-22.
- [2] T.G. Bhaskar and V. Lakshmikantham, Fixed point theorems in partially ordered metric spaces and applications, Nonlinear Anal. 65(2006), 1379-1393.
- [3] D.W. Boyd and T.S.W. Wong, On nonlinear contractions, Proc. Amer. Math. Soc. 20(1969), 458-464.
- [4] S. Chandok and T.D. Narang, Some common fixed point theorems for Banach operator pairs with applications in best approximation, Nonlinear Analysis-TMA 73(2010), 105-109.
- [5] S. Chandok and T.D. Narang, Common fixed points and invariant approximation for Gregus type contraction mappings, Rendiconti Circolo Mat. Palermo 60(2011), 203-214.
- [6] S. Chandok and T.D. Narang, Common fixed points and invariant approximation for Banach operator pairs with Ćirić type nonexpansive mappings, Hacettepe J. Math Stat. 40(2011), 871-883.
- [7] S. Chandok, Some common fixed point theorems for generalized f -weakly contractive mappings, J. Appl. Math. Informatics 29(2011), 257-265.
- [8] S. Chandok, Some common fixed point theorems for generalized nonlinear contractive mappings, Comput. Math. Appl. 62(2011), 3692-3699.
- [9] S. Chandok, Some common fixed point theorems for Ćirić type contractive mappings, Tamkang J. Math. 43 (2) (2012), 187-202.
- [10] S. Chandok, Common fixed points, invariant approximation and generalized weak contractions, Internat. J. Math. Math. Sci. 2012 Article ID 102980.
- [11] S. Chandok, Common fixed points for generalized nonlinear contractive mappings in metric spaces, Mat. Vesnik(in press).
- [12] S. Chandok, J. Liang and D. O'Regan, Common fixed points and invariant approximations for non-commuting contraction mappings in strongly convex metric spaces, J. Nonlinear Convex Analysis Vol 14 (2013).
- [13] B.S. Choudhury, Unique fixed point theorem for weakly C -contractive mappings, Kathmandu University J. Sci. Engg. Tech. 5(2009), 6-13.
- [14] L. Ćirić, N. Hussain and N. Ćakić, Common fixed points for Ćirić type f -contraction with applications, Publ. Math. Debrecen 4317(2009), 1-19.
- [15] J. Harjani, B. López and K. Sadarangani, Fixed point theorems for weakly C -contractive mapping in ordered metric spaces, Computers and Mathematics with Applications 61(4)(2011), 790-796.
- [16] M.S. Khan, M. Swaleh and S. Sessa, Fixed point theorems by altering distances between the points, Bull. Aust. Math. Soc. 30(1984), 1-9.
- [17] J.J. Nieto and R.R. López, Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations, Order 22(2005), 223-239.
- [18] A.C.M. Ran and M.C.B. Reurings, A fixed point theorem in partially ordered sets and some applications to matrix equations, Proc. Amer. Math. Soc. 132(5)(2004), 1435-1443.

- [19] S. Reich, Some fixed point problems, *Atti Acad. Naz. Lincei Ren. Cl. Sci. Fis. Mat. Natur.* 57(1975), 194-198.
- [20] B.E. Rhoades, Some theorems on weakly contractive maps, *Nonlinear Anal* 47(2001), 2683-2693.