



## HARDY-ROGERS TYPE MAPPINGS ON DISLOCATED QUASI METRIC SPACES

ANANTACHAI PADCHAROEN<sup>\*1</sup>, DHANANJAY GOPAL<sup>2</sup> AND  
NAKNIMIT AKKASRIWORN<sup>1</sup>

<sup>1</sup> Department of Mathematics, Faculty of Science and Technology, Rambhai Barni Rajabhat University, Chanthaburi 22000, Thailand

<sup>2</sup> Department Of Applied Mathematics & Humanities, Sardar Vallabhbhai National Institute of Technology, Surat, 395-007, Gujarat, India

---

**ABSTRACT.** In this paper, We prove some common fixed point results for two  $\alpha$ -dominated mappings satisfying Hardy-Rogers Type on a closed ball of left (right)  $K$ -sequentially complete dislocated quasi-metric space and give some example for support our result.

**KEYWORDS:** Hardy-Rogers Type, Dislocated Quasi Metric Spaces, Quasi Metric Spaces.

**AMS Subject Classification:** 47H09, 47H10, 49M05

---

### 1. INTRODUCTION

The partial metric spaces have applications in theoretical computer science (see [14]). The notion of dislocated topologies has useful applications in the context of logic programming semantics (see [15]). Dislocated metric (metric-like) spaces (see [4, 16, 17, 18]) are generalizations of partial metric spaces. Furthermore, dislocated quasi metric spaces (quasi-metric-like spaces) generalize the idea of dislocated metric spaces and quasi-partial metric spaces.

Samet et al. [19] introduced the notion of  $\alpha$ -admissible mappings. They weakened and generalized the contractive condition and several other known results.

In this paper, we proof common fixed point results for two  $\alpha$ -dominated mappings in a closed ball in complete dislocated quasi metric space, under Hardy-Rogers Type.

### 2. PRELIMINARIES

**Definition 2.1.** [10] Let  $X$  be a nonempty set. A quasi-partial metric on  $X$  is a function  $q : X \times X \rightarrow \mathbb{R}^+$  satisfying, for all  $x, y, z \in X$ ,

---

<sup>\*</sup> Corresponding author.

Email address : apadcharoen@yahoo.com, gopal.dhananjay@rediffmail.com, boyjuntaburi@hotmail.com.

- (a)  $0 \leq q(x, x) = q(x, y) = q(y, y)$  implies  $x = y$  (equality),
- (b)  $q(x, x) \leq q(y, x)$  (small self-distances),
- (c)  $q(x, x) \leq q(x, y)$  (small self-distances),
- (d)  $q(x, y) + q(z, z) \leq q(x, z) + q(z, y)$  (triangle inequality).

The pair  $(X, q)$  is called a quasi-partial metric space.

**Definition 2.2.** [13] Let  $X$  be a nonempty set. A function  $d_q : X \times X \rightarrow [0, \infty)$  is called a dislocated quasi metric (or simply  $d_q$ -metric) if the following conditions hold for any  $x, y, z \in X$  :

- (a) If  $d_q(x, y) = d_q(y, x) = 0$ , then  $x = y$ ,
- (b)  $d_q(x, y) \leq d_q(x, z) + d_q(z, y)$ .

In this case, the pair  $(X, d_q)$  is called a dislocated quasi metric space.

It is clear that, if  $d_q(x, y) = d_q(y, x) = 0$ , then from (a) we have  $x = y$ . But, if  $x = y$ , then  $d_q(x, y)$  may not be 0. It can be observed that, if  $d_q(x, y) = d_q(y, x)$  for all  $x, y \in X$ , then  $(X, d_q)$  becomes a dislocated metric space (metric-like space)[1, 4, 5, 6, 9]. We will denote by  $(X, d_l)$  a dislocated metric space. For  $x \in X$  and  $\epsilon > 0$ ,  $B_{d_q}(x, \epsilon) = \{y \in X : d_q(x, y) \leq \epsilon\}$  is a closed ball in  $(X, d_q)$ . Every quasi-partial metric space is a dislocated quasi metric space, but the converse is not true in general.

**Example 2.3.** If  $X = \mathbb{R}^+ \cup \{0\}$ , then  $d_q(x, y) = x + \max\{x, y\}$  defines a dislocated quasi metric  $d_q$  on  $X$ . But, it is not a quasi-partial metric space. Indeed,

$$d_q(3, 3) = 6 > d_q(2, 3) = 5.$$

Reilly et al. [11] introduced the notion of left (right)  $K$ -Cauchy sequence and left (right)  $K$ -sequentially complete spaces.

**Definition 2.4.** Let  $(X, d_q)$  be a dislocated quasi metric space.

- (a) A sequence  $\{x_n\}$  in  $(X, d_q)$  is called left(right)  $K$ -Cauchy if  $\forall \epsilon > 0, \exists n_0 \in \mathbb{N}$  such that  $\forall n > m \leq n_0, d_q(x_n, x_m) < \epsilon$  (respectively  $d_q(x_n, x_m) < \epsilon$ ).
- (b) A sequence  $\{x_n\}$  in  $(X, d_q)$  dislocated quasi-converges (for short  $d_q$ -converges) to  $x$  if  $\lim_{n \rightarrow \infty} d_q(x_n, x) = \lim_{n \rightarrow \infty} d_q(x, x_n) = 0$ . In this case, the point  $x$  is called a  $d_q$ -limit of  $\{x_n\}$ .
- (c)  $(X, d_q)$  is called left (right)  $K$ -sequentially complete if every left (right)  $K$ -Cauchy sequence in  $(X, d_q)$ ,  $d_q$  -converges to a point  $x \in X$  such that  $d_q(x, x) = 0$ .

One can easily observe that every complete dislocated quasi metric space is also left  $K$ -sequentially complete dislocated quasi metric space, but the converse is not true in general.

**Remark 2.5.** [3] It is easy to see that, if  $x_n \in B_{d_q}(x_0, r)$  for all  $n \in \mathbb{N}$  and for some  $x_0 \in X, r > 0$ , and the sequence  $\{x_n\}$ ,  $d_q$ -converges to a point  $z \in X$ , then  $z \in B_{d_q}(x_0, r)$ .

**Definition 2.6.** [12] Let  $(X, q)$  be a quasi-partial metric space.

- (a) A sequence  $\{x_n\}$  in  $(X, q)$  is called 0-Cauchy if  $\lim_{n, m \rightarrow \infty} q(x_n, x_m) = 0$  or  $\lim_{n, m \rightarrow \infty} q(x_m, x_n) = 0$ .
- (b) The space  $(X, q)$  is called 0-complete if every 0-Cauchy sequence in  $X$  converges to a point  $x \in X$  such that  $q(x, x) = 0$ .

**Remark 2.7.** [3] By definitions, one can easily observe that if  $X$  is a 0-complete quasi-partial metric space then it is also a  $K$ -sequentially complete dislocated quasi metric space. But a  $K$ -sequentially complete dislocated quasi metric space may not be a 0-complete quasi-partial metric space. Therefore, the results in a  $K$ -sequentially complete dislocated quasi metric space are more general than those in a 0-complete quasi-partial metric space.

Let  $X$  be a non-empty set and  $T, f : X \rightarrow X$  be two mappings. A point  $y \in X$  is called a point of coincidence of  $T$  and  $f$  if there exists a point  $x \in X$  such that  $y = Tx = fx$ , here  $x$  is called a coincidence point of  $T$  and  $f$ . The mappings  $T, f$  are said to be weakly compatible if they commute at their coincidence points i.e.,  $Tfx = fTx$  whenever  $Tx = fx$ .

Let  $\Psi$  denote the family of all nondecreasing functions  $\psi : [0, +\infty) \rightarrow [0, +\infty)$  such that  $\sum_{n=1}^{\infty} \psi^n(t) < +\infty$  for all  $t \geq 0$ , where  $\psi^n$  is the  $n^{\text{th}}$  iterate of  $\psi$ . The following lemma is a consequence of definition of  $\Psi$ .

**Lemma 2.8.** *If  $\psi \in \Psi$ , then  $\psi(t) < t$  for all  $t > 0$ .*

**Definition 2.9.** [3] Let  $(X, d_q)$  be a dislocated quasi metric space,  $A \subseteq X$ ,  $T : X \rightarrow X$  be a selfmapping and  $\alpha : X \times X \rightarrow [0, +\infty)$ . Then:

- (a) The mapping  $T$  is said to be  $\alpha$ -dominated on  $A$ , if  $\alpha(x, Tx) \geq 1$  for all  $x \in A$ .
- (b) The function  $\alpha$  is said to be a triangular function on  $A$ , if  $\alpha(x, y) \geq 1$  and  $\alpha(y, z) \geq 1$  implies that  $\alpha(x, z) \geq 1$  for all  $x, y, z \in A$ .
- (b)  $(X, d_q)$  is  $\alpha$ -regular on  $A$  if for any sequence  $\{x_n\}$  in  $A$  such that  $\alpha(x_n, x_{n+1}) \geq 1$  for all  $n \geq 0$  and  $x_n \rightarrow x \in A$  as  $n \rightarrow \infty$  we have  $\alpha(x_n, x) \geq 1$  for all  $n \geq 0$ .

It is clear that if  $T$  is an  $\alpha$ -dominated mapping on  $X$  then  $T$  is  $\alpha$ -dominated on each subset of  $X$ , but  $T$  can be  $\alpha$ -dominated on some  $A \subseteq X$ , without being  $\alpha$ -dominated mapping on  $X$ .

### 3. MAIN RESULTS

**Theorem 3.1.** *Let  $(X, d_q)$  be a left  $K$ -sequentially complete dislocated quasi metric space and  $T, S : X \rightarrow X$  be two mappings. Let  $x_0 \in X$ ,  $r > 0$  and there exists a function  $\alpha : X \times X \rightarrow [0, +\infty)$  such that  $S$  and  $T$  are  $\alpha$ -dominated mappings on  $\overline{B_{d_q}(x_0, r)}$ . Suppose that  $x_0 \in B_{d_q}(x_0, r)$  and there exist nonnegative real numbers  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + 2\delta \in (0, 1)$  and the following condition holds: if  $\alpha(x, y) \geq 1$  or  $\alpha(y, x) \geq 1$  and  $x, y \in \overline{B_{d_q}(x_0, r)}$ , then*

$$d_q(Sx, Ty) \leq \beta d_q(x, y) + \gamma[d_q(x, Sx) + d_q(y, Ty)] + \delta[d_q(y, Sx) + d_q(x, Ty)] \quad (3.1)$$

$$d_q(Tx, Sy) \leq \beta d_q(x, y) + \gamma[d_q(x, Tx) + d_q(y, Sy)] + \delta[d_q(y, Tx) + d_q(x, Sy)] \quad (3.2)$$

and

$$d_q(x_0, Sx_0) \leq (1 - \lambda)r, \quad (3.3)$$

where  $\lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta}$ . Suppose that  $(X, d_q)$  is  $\alpha$ -regular on  $\overline{B_{d_q}(x_0, r)}$ . Then there exists a common fixed point  $z \in \overline{B_{d_q}(x_0, r)}$  of  $S$  and  $T$ . Moreover,  $d_q(z, z) = 0$ .

*Proof.* Let  $x_0 \in X$ , define  $x_1 = Sx_0$  and  $x_2 = Tx_1$ . Continuing this process, we construct a sequence  $\{x_n\}$  of points in  $X$ , such that

$$x_{2k+1} = Sx_{2k} \text{ and } x_{2k+2} = Tx_{2k+1}, \quad \forall k = 0, 1, 2, \dots$$

By mathematical induction, we can show that

$$\begin{cases} x_{n+1} \in \overline{B_{d_q}(x_0, r)}, \alpha(x_n, x_{n+1}) \geq 1, \\ d_q(x_n, x_{n+1}) \leq \lambda^n d_q(x_0, x_1), \quad \forall n \in \mathbb{N}. \end{cases} \quad (P_n)$$

By using (3.3) and  $0 < \lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta} < 1$ , we obtain

$$d_q(x_0, x_1) = d_q(x_0, Sx_0) \leq (1 - \lambda)r \leq r.$$

Hence,  $x_1 \in \overline{B_{d_q}(x_0, r)}$ . Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(x_0, Sx_0) = \alpha(x_0, x_1) \geq 1$ . Therefore, using (3.1), we get that

$$\begin{aligned} d_q(x_1, x_2) &= d_q(Sx_0, Tx_1) \\ &\leq \beta d_q(x_0, x_1) + \gamma [d_q(x_0, Sx_0) + d_q(x_1, Tx_1)] \\ &\quad + \delta [d_q(x_1, Sx_0) + d_q(x_0, Tx_1)] \\ &= \beta d_q(x_0, x_1) + \gamma [d_q(x_0, x_1) + d_q(x_1, x_2)] \\ &\quad + \delta [d_q(x_1, x_1) + d_q(x_0, x_2)] \\ &\leq \beta d_q(x_0, x_1) + \gamma [d_q(x_0, x_1) + d_q(x_1, x_2)] \\ &\quad + \delta [d_q(x_0, x_1) + d_q(x_1, x_2)] \end{aligned}$$

Thus,

$$d_q(x_1, x_2) \leq \lambda d_q(x_0, x_1). \quad (3.4)$$

By using (3.4), we get that

$$\begin{aligned} d_q(x_0, x_2) &\leq d_q(x_0, x_1) + d_q(x_1, x_2) \leq d_q(x_0, x_1) + \lambda d_q(x_0, x_1) \\ &= (1 + \lambda) d_q(x_0, x_1) \leq (1 + \lambda)(1 - \lambda)r = (1 - \lambda^2)r \leq r. \end{aligned}$$

Hence,  $x_2 \in \overline{B_{d_q}(x_0, r)}$ . Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(x_0, Sx_0) = \alpha(x_1, Tx_1) \geq 1$ . Therefore, from (3.1) holds and using (3.2), we get that

$$\begin{aligned} d_q(x_2, x_3) &= d_q(Tx_1, Sx_2) \\ &\leq \beta d_q(x_1, x_2) + \gamma [d_q(x_1, Tx_1) + d_q(x_2, Sx_2)] \\ &\quad + \delta [d_q(x_2, Tx_1) + d_q(x_1, Sx_2)] \\ &= \beta d_q(x_1, x_2) + \gamma [d_q(x_1, x_2) + d_q(x_2, x_3)] \\ &\quad + \delta [d_q(x_2, x_2) + d_q(x_1, x_3)] \\ &\leq \beta d_q(x_1, x_2) + \gamma [d_q(x_1, x_2) + d_q(x_2, x_3)] \\ &\quad + \delta [d_q(x_1, x_2) + d_q(x_2, x_3)] \end{aligned}$$

By using (3.4), we get that

$$d_q(x_2, x_3) \leq \lambda d_q(x_1, x_2) \leq \lambda^2 d_q(x_0, x_1). \quad (3.5)$$

It follows from (3.4) and (3.5) that

$$\begin{aligned} d_q(x_0, x_3) &\leq d_q(x_0, x_1) + d_q(x_1, x_2) + d_q(x_2, x_3) \\ &\leq d_q(x_0, x_1) + \lambda d_q(x_0, x_1) + \lambda^2 d_q(x_0, x_1) \\ &= (1 + \lambda + \lambda^2) d_q(x_0, x_1) = \frac{1 - \lambda^3}{1 - \lambda} d_q(x_0, x_1) \\ &\leq \frac{1 - \lambda^3}{1 - \lambda} (1 - \lambda)r = (1 - \lambda^3)r \leq r. \end{aligned}$$

Hence,  $x_3 \in \overline{B_{d_q}(x_0, r)}$ . Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(x_0, Sx_0) = \alpha(x_1, Tx_1) \geq 1$ . Therefore, from (3.1) holds. Suppose,  $(P_1), (P_2), \dots, (P_i)$

be the inductive hypothesis. We shall show that  $(P_{i+1})$  holds. For this, we consider two possible cases. First, suppose that  $i$  is even. Then, since  $\alpha(x_i, x_{i+1}) \geq 1$  and using (3.1), we get that

$$\begin{aligned} d_q(x_{i+1}, x_{i+2}) &= d_q(Sx_i, Tx_{i+1}) \\ &\leq \beta d_q(x_i, x_{i+1}) + \gamma[d_q(x_i, Sx_i) + d_q(x_{i+1}, Tx_{i+1})] \\ &\quad + \delta[d_q(x_{i+1}, Sx_i) + d_q(x_i, Tx_{i+1})] \\ &= \beta d_q(x_i, x_{i+1}) + \gamma[d_q(x_i, x_{i+1}) + d_q(x_i, x_{i+2})] \\ &\quad + \delta[d_q(x_{i+1}, x_{i+1}) + d_q(x_i, x_{i+2})] \\ &\leq \beta d_q(x_i, x_{i+1}) + \gamma[d_q(x_i, x_{i+1}) + d_q(x_{i+1}, x_{i+2})] \\ &\quad + \delta[d_q(x_i, x_{i+1}) + d_q(x_{i+1}, x_{i+2})] \end{aligned}$$

Since  $(P_i)$  holds, we get that

$$d_q(x_{i+1}, x_{i+2}) \leq \lambda d_q(x_i, x_{i+1}) \leq \lambda^{i+1} d_q(x_0, x_1).$$

Thus,

$$\begin{aligned} d_q(x_0, x_{i+2}) &\leq d_q(x_0, x_1) + d_q(x_1, x_2) + \cdots + d_q(x_{i+1}, x_{i+2}) \\ &\leq (1 + \lambda + \lambda^2 + \cdots + \lambda^{i+2}) d_q(x_0, x_1) \\ &\leq \frac{1 - \lambda^{i+2}}{1 - \lambda} (1 - \lambda) r \leq (1 - \lambda^{i+2}) r \leq r. \end{aligned}$$

Hence,  $x_{i+2} \in \overline{B_{d_q}(x_0, r)}$ . Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(x_{i+1}, Sx_{i+1}) = \alpha(x_{i+1}, x_{i+2}) \geq 1$ . Therefore,  $(P_{i+1})$  holds. Similarly, one can see that if  $i$  is odd, then  $(P_{i+1})$  holds, which completes the inductive proof. Thus, we can write

$$d_q(x_n, x_{n+1}) \leq \lambda^n d_q(x_0, x_1), \quad \forall n \in \mathbb{N}. \quad (3.6)$$

Next, we will show that the sequence  $\{x_n\}$  is a left  $K$ -Cauchy sequence. Indeed, for  $n, m \in \mathbb{N}$  with  $m > n$  using (3.7) we have

$$\begin{aligned} d_q(x_n, x_m) &\leq d_q(x_n, x_{n+1}) + d_q(x_{n+1}, x_{n+2}) + \cdots + d_q(x_{m-1}, x_m) \\ &\leq \lambda^n d_q(x_0, x_1) + \lambda^{n+1} d_q(x_0, x_1) + \cdots + \lambda^{m-1} d_q(x_0, x_1). \end{aligned}$$

Thus,

$$d_q(x_n, x_m) \leq \frac{\lambda^n}{1 - \lambda} d_q(x_0, x_1), \quad \forall n, m \in \mathbb{N}, m > n. \quad (3.7)$$

Since  $0 < \lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta} < 1$ , for every  $\epsilon > 0$ , we can choose  $n_0 \in \mathbb{N}$  such that  $\lambda^n < \frac{1 - \lambda}{d_q(x_0, x_1)} \epsilon$  for all  $n > n_0$ . Therefore, it follows from (3.7) that

$$d_q(x_n, x_m) < \epsilon, \quad \forall m > n > n_0.$$

Therefore, the sequence  $\{x_n\}$  is a left  $K$ -Cauchy sequence in  $X$ . By left  $K$ -sequential completeness of  $X$ , there exists  $x \in X$  such that

$$\lim_{n \rightarrow \infty} d_q(x_n, z) = \lim_{n \rightarrow \infty} d_q(z, x_n) = 0. \quad (3.8)$$

We will show that  $z$  is a common fixed point of the mappings  $S$  and  $T$ . By Remark 2.5, we have  $z \in B_{d_q}(x_0, r)$ . Now, by the assumption we have for all  $n \in \mathbb{N}$ , therefore

for any  $n \in \mathbb{N}$ , we have

$$\begin{aligned}
 d_q(z, Sz) &\leq d_q(z, x_{2n+2}) + d_q(x_{2n+2}, Sz) \\
 &\leq d_q(z, x_{2n+2}) + d_q(Tx_{2n+1}, Sz) \\
 &\leq d_q(z, x_{2n+2}) + \beta d_q(x_{2n+1}, z) \\
 &\quad + \gamma [d_q(x_{2n+1}, Sx_{n+1}) + d_q(z, Sz)] \\
 &\quad + \delta [d_q(z, Tx_{2n+1}) + d_q(x_{2n+1}, Sz)] \\
 &\leq d_q(z, x_{2n+2}) + \beta d_q(x_{2n+1}, z) \\
 &\quad + \gamma [d_q(x_{2n+1}, Sx_{n+1}) + d_q(z, Sz)] \\
 &\quad + \delta [d_q(z, x_{2n+2}) + d_q(x_{2n+1}, z) + d_q(z, Sz)].
 \end{aligned}$$

By using (3.7) and (3.8), we obtain

$$(1 - \gamma + \delta)d_q(z, Sz) \leq 0 \quad (3.9)$$

which implies that  $d_q(z, Sz) = 0$ . Similarly, one can show that  $d_q(Sz, z) = 0$ . Thus,  $d_q(z, Sz) = d_q(Sz, z) = 0$ , i.e.,  $z = Sz$ . Similarly, one can show that  $z = Tz$ .

Hence,  $S$  and  $T$  have a common fixed point  $z \in \overline{B_{d_q}(x_0, r)}$ . As is an dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(z, Sz) = \alpha(z, z) \geq 1$ . Therefore,

$$\begin{aligned}
 d_q(z, z) &\leq d_q(Sz, Tz) \\
 &\leq \beta d_q(z, z) + \gamma [d_q(z, Sz) + d_q(z, Tz)] \\
 &\quad + \delta [d_q(z, Sz) + d_q(z, Tz)] \\
 &\leq (\beta + 2\gamma + 2\delta)d_q(z, z),
 \end{aligned}$$

and this implies that

$$d_q(z, z) = 0.$$

□

**Example 3.2.** Let  $X = \mathbb{Q}^+ \cup \{0\}$  and let  $d_q : X^2 \times X^2 \rightarrow X$  be defined by  $d_q((x_1, y_1), (x_2, y_2)) = x_1 + 4y_1 + \frac{x_2}{4} + y_2$ . Then it is easy to show that  $(X^2, d_q)$  is a left  $K$ -sequentially complete dislocated quasi metric space. If  $(x_0, y_0) = (4, 1)$ ,  $r = 28$ , then

$$\overline{B_{d_q}((4, 1), 28)} = \{(x, y) \in X : x + 4y \leq 42\}.$$

In particular,  $(4, 1) \in \overline{B_{d_q}((4, 1), 28)}$ .

Let  $S, T : X^2 \rightarrow X^2$  be defined by

$$S(x, y) = \begin{cases} \left(\frac{x}{7}, \frac{y}{7}\right), & \text{if } x + 4y \leq 42 \\ (2x^2 - 2, 4x + 5), & \text{if } x + 4y > 42 \end{cases}$$

and

$$T(x, y) = \begin{cases} \left(\frac{x}{6}, \frac{y}{9}\right), & \text{if } x + 4y \leq 42 \\ (3x^2 - 3, y), & \text{if } x + 4y > 42. \end{cases}$$

Also, define  $\alpha : X^2 \times X^2 \rightarrow [0, \infty)$  by

$$\alpha((x_1, y_1), (x_2, y_2)) = \begin{cases} 1, & \text{if } \frac{x_1}{4} + y_1 + x_2 + y_2 \leq 42 \\ 0, & \text{if } \frac{x_1}{4} + y_1 + x_2 + y_2 > 42. \end{cases}$$

Clearly,  $S$  and  $T$  are  $\alpha$ -dominated mappings on  $\overline{B_{d_q}((4, 1), 28)}$ . Let  $\beta = \frac{1}{7}$ ,  $\gamma = \delta = \frac{1}{10}$ , then  $\lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta} = \frac{3}{10} \in (0, 1)$  and  $(1 - \lambda)r = 16$ ,  $d_q((x_0, y_0), S(x_0, y_0)) = d_q((4, 1), S(4, 1)) = \frac{104}{7} < 16 = (1 - \lambda)r$ . Observe that, for  $(43, 0) \notin \overline{B_{d_q}((4, 1), 28)}$ , we have  $d_q(S(43, 0), T(43, 0)) = d_q((3696, 5), (5544, 0)) = 5104$ ,  $d_q((43, 0), T(43, 0)) + d_q((43, 0), S(43, 0)) = 2401$ , and  $d_q((43, 0), (43, 0)) = \frac{215}{4}$ . Hence, there are no  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + \delta \in (0, 1)$  and (3.1) is satisfied. So the contractive condition does not hold on  $X^2$ . On the other hand, if  $(x_1, y_1), (x_2, y_2) \in \overline{B_{d_q}((4, 1), 28)}$ , then

$$\begin{aligned} d_q(S(x_1, y_1), T(x_2, y_2)) &= d_q\left(\left(\frac{x_1}{7}, \frac{y_1}{7}\right), \left(\frac{x_2}{6}, \frac{y_2}{9}\right)\right) \\ &= \frac{x_1}{7} + \frac{4y_1}{7} + \frac{x_2}{24} + \frac{y_2}{9} \\ &\leq \frac{1}{7}d_q((x_1, y_1), (x_2, y_2)) \\ &\quad + \frac{1}{10}[d_q((x_1, y_1), S(x_1, y_1)) + d_q((x_2, y_2), T(x_2, y_2))] \\ &\quad + \frac{1}{10}[d_q((x_2, y_2), S(x_1, y_1)) + d_q((x_1, y_1), T(x_2, y_2))]. \end{aligned}$$

Also,

$$\begin{aligned} d_q(T(x_1, y_1), S(x_2, y_2)) &= d_q\left(\left(\frac{x_1}{6}, \frac{y_1}{9}\right), \left(\frac{x_2}{7}, \frac{y_2}{7}\right)\right) \\ &= \frac{x_1}{6} + \frac{4y_1}{9} + \frac{x_2}{28} + \frac{y_2}{7} \\ &\leq \frac{1}{7}d_q((x_1, y_1), (x_2, y_2)) \\ &\quad + \frac{1}{10}[d_q((x_1, y_1), S(x_1, y_1)) + d_q((x_2, y_2), T(x_2, y_2))] \\ &\quad + \frac{1}{10}[d_q((x_2, y_2), S(x_1, y_1)) + d_q((x_1, y_1), T(x_2, y_2))]. \end{aligned}$$

Therefore, all the conditions of Theorem 3.1 are satisfied. Moreover,  $(0, 0)$  is the common fixed point of  $S$  and  $T$ .

**Corollary 3.3.** *Let  $(X, d_q)$  be a left  $K$ -sequentially complete dislocated quasi metric space and  $S : X \rightarrow X$  be a mapping. Let  $x_0 \in X$ ,  $r > 0$  and there exists a function  $\alpha : X \times X \rightarrow [0, +\infty)$  such that  $S$  be an  $\alpha$ -dominated mappings on  $B_{d_q}(x_0, r)$ . Suppose that  $x_0 \in B_{d_q}(x_0, r)$  and there exist nonnegative real numbers  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + \delta \in (0, 1)$  and the following condition holds: if  $\alpha(x, y) \geq 1$  or  $\alpha(y, x) \geq 1$  and  $x, y \in B_{d_q}(x_0, r)$ , then*

$$d_q(Sx, Sy) \leq \beta d_q(x, y) + \gamma [d_q(x, Sx) + d_q(y, Sy)] + \delta [d_q(y, Sx) + d_q(x, Sy)]$$

and

$$d_q(x_0, Sx_0) \leq (1 - \lambda)r,$$

where  $\lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta}$ . Suppose that  $(X, d_q)$  is  $\alpha$ -regular on  $\overline{B_{d_q}(x_0, r)}$ . Then there exists a point  $z \in \overline{B_{d_q}(x_0, r)}$  such that  $z = Sz$  and  $d_q(z, z) = 0$ .

*Proof.* Letting  $T = S$  in Theorem 3.1, we obtain the following result.  $\square$

**Corollary 3.4.** *Let  $(X, d)$  be a complete dislocated metric space and  $S, T : X \rightarrow X$  be two mappings. Let  $x_0 \in X$ ,  $r > 0$  and there exists a function  $\alpha : X \times X \rightarrow [0, +\infty)$  such that  $S$  and  $T$  are  $\alpha$ -dominated mappings on  $\overline{B_d(x_0, r)}$ . Suppose that  $x_0 \in \overline{B_d(x_0, r)}$  and there exist nonnegative real numbers  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + \delta \in (0, 1)$  and the following condition holds: if  $\alpha(x, y) \geq 1$  or  $\alpha(y, x) \geq 1$  and  $x, y \in B_d(x_0, r)$ , then*

$$d(Sx, Ty) \leq \beta d(x, y) + \gamma [d(x, Sx) + d(y, Ty)] + \delta [d(y, Sx) + d(x, Ty)]$$

and

$$d(x_0, Sx_0) \leq (1 - \lambda)r,$$

where  $\lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta}$ . Suppose that  $(X, d)$  is  $\alpha$ -regular on  $\overline{B_d(x_0, r)}$ . Then there exists a point  $z \in B_d(x_0, r)$  such that  $z = Sz$  and  $d(z, z) = 0$ .

*Proof.* By Theorem 3.1, we obtain the following result.  $\square$

**Theorem 3.5.** *Suppose that all the conditions of Theorem 3.1 are satisfied. In addition suppose that:*

- (a) *The function  $\alpha$  is a triangular function on  $\overline{B_{d_q}(x_0, r)}$ .*
- (b) *For  $x, y \in \overline{B_{d_q}(x_0, r)}$  there exists  $u_0 \in \overline{B_{d_q}(x_0, r)}$  such that  $\alpha(x, u_0) \geq 1, \alpha(y, u_0) \geq 1$ .*
- (c) *For all  $u \in \overline{B_{d_q}(x_0, r)}$  such that  $\alpha(Sx_0, u) \geq 1$  the following condition holds*  
 $d_q(x_0, Sx_0) + d_q(u, Tu) + d_q(u, Sx_0) + d_q(x_0, Tu) \leq d_q(x_0, u) + d_q(Sx_0, Tu)$ .

Then  $S$  and  $T$  have a unique common fixed point  $z \in \overline{B_{d_q}(x_0, r)}$  and  $d_q(z, z) = 0$ .

*Proof.* Define the sequence  $\text{fxng}$  as in the proof Theorem 3.1. Then,  $\{x_n\}, d_q$ -converges to a common fixed point  $z \in \overline{B_{d_q}(x_0, r)}$  of the mappings  $S$  and  $T$  such that  $\alpha(x_n, z) \geq 1$  for all  $n \geq 0$ ,  $(P_n)$  holds and  $d_q(z, z) = 0$ . In order to prove uniqueness of  $z$ , suppose that  $z^*$  is another point in  $\overline{B_{d_q}(x_0, r)}$  such that  $z^* = Sz^* = Tz^*$ . Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(z^*, Sz^*) = \alpha(z^*, z^*) \geq 1$ . Therefore,

$$\begin{aligned} d_q(z^*, z^*) &\leq d_q(Sz^*, Tz^*) \\ &\leq \beta d_q(z^*, z^*) + \gamma [d_q(z^*, Sz^*) + d_q(z^*, Tz^*)] \\ &\quad + \delta [d_q(z^*, Sz^*) + d_q(z^*, Tz^*)] \\ &\leq (\beta + 2\gamma + 2\delta) d_q(z^*, z^*), \end{aligned}$$

and this implies that

$$d_q(z^*, z^*) = 0.$$

By assumption, there exists a point  $u_0 \in \overline{B_{d_q}(x_0, r)}$  such that  $\alpha(z, u_0) \geq 1$  and  $\alpha(z^*, u_0) \geq 1$ . Define a sequence  $\{u_n\}$  in  $X$  such that,

$$u_{2k+1} = Su_{2k} \text{ and } u_{2k+2} = Tu_{2k+1}, \quad \forall k = 0, 1, 2, \dots$$

By mathematical induction, we can show that

$$\left\{ \begin{array}{l} \alpha(u_n, u_{n+1}) \geq 1, \alpha(x_n, u_n) \geq 1, \quad \forall n \in \mathbb{N}; \\ d_q(u_n, u_{n+1}) \leq \lambda^n d_q(u_0, u_1), \quad \forall n \in \mathbb{N}; \\ d_q(x_n, z_n) \leq \lambda^n r, u_n \in \overline{B_{d_q}(x_0, r)}, \quad \forall n \in \mathbb{N}. \end{array} \right. \quad (P'_n)$$

Since  $T$  is  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(u_0, Tu_0) = \alpha(u_0, u_1) \geq 1$ . Since  $\alpha$  is triangular function on  $\overline{B_{d_q}(x_0, r)}$ , and  $\alpha(x_n, z) \geq 1, \alpha(z, u_0) \geq 1$ , we have  $\alpha(x_n, u_0) \geq 1$  for all  $n \geq 0$ . Therefore, using (c), we get that

$$\begin{aligned} d_q(x_1, u_1) &= d_q(Sx_0, Tu_0) \\ &\leq \beta d_q(x_0, u_0) + \gamma[d_q(x_0, Sx_0) + d_q(u_0, Tu_0)] \\ &\quad + \delta[d_q(u_0, Sx_0) + d_q(x_0, Tu_0)] \\ &\leq \beta d_q(x_0, u_0) + \gamma[d_q(x_0, u_0) + d_q(Sx_0, Tu_0)] \\ &\quad + \delta[d_q(x_0, u_0) + d_q(Sx_0, Tu_0)] \\ &\leq \beta d_q(x_0, u_0) + \gamma[d_q(x_0, u_0) + d_q(x_1, u_1)] \\ &\quad + \delta[d_q(u_0, x_0) + d_q(x_1, u_1)]. \end{aligned}$$

Thus,

$$d_q(x_1, u_1) \leq \frac{\beta + \gamma + \delta}{1 - \gamma - \delta} d_q(x_0, u_0) = \lambda d_q(x_0, u_0) \leq \lambda r. \quad (3.10)$$

Since  $u_0 \in \overline{B_{d_q}(x_0, r)}$ , using (3.10), we get

$$\begin{aligned} d_q(x_0, u_1) &\leq d_q(x_0, x_1) + d_q(x_1, u_1) \\ &\leq (1 - \lambda)r + \lambda d_q(x_0, u_0) \\ &\leq (1 - \lambda)r + \lambda r \leq r \end{aligned}$$

Hence,  $u_1 \in \overline{B_{d_q}(x_0, r)}$ . Since  $\alpha(u_0, u_1) \geq 1$ , by using (3.2), we get that

$$d_q(u_1, u_2) \leq \frac{\beta + \gamma + \delta}{1 - \gamma - \delta} d_q(u_0, u_1) = \lambda d_q(u_0, u_1).$$

Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(u_1, Su_1) = \alpha(u_1, u_1) \geq 1$ . As,  $\alpha$  is a triangular function on  $\overline{B_{d_q}(x_0, r)}$ , and  $\alpha(x_1, u_0) \geq 1, \alpha(u_0, u_1) \geq 1$ , we have  $\alpha(x_1, u_1) \geq 1$ . Therefore, from  $(P'_1)$  holds. Since  $\alpha(u_1, u_2) \geq 1$  and using (3.1), we get that

$$d_q(u_2, u_3) \leq \lambda d_q(u_1, u_2) \leq \lambda^2 d_q(u_0, u_1).$$

Since  $\alpha(x_1, u_1) \geq 1$ , using (3.2) that

$$\begin{aligned} d_q(x_2, u_2) &= d_q(Tx_1, Su_1) \\ &\leq \beta d_q(x_1, u_1) + \gamma[d_q(x_1, Tx_1) + d_q(u_1, Su_1)] \\ &\quad + \delta[d_q(u_1, Tx_1) + d_q(x_1, Su_1)] \\ &\leq \beta d_q(x_1, x_2) + \gamma\lambda[d_q(x_0, Sx_0) + d_q(u_0, Tu_0)] \\ &\quad + \delta\lambda[d_q(u_0, Sx_0) + d_q(x_0, Tu_0)] \end{aligned}$$

which gives with (c)

$$\begin{aligned} d_q(x_2, u_2) &\leq \beta d_q(x_1, x_2) + \gamma\lambda[d_q(x_0, u_0) + d_q(Sx_0, Tu_0)] \\ &\quad + \delta\lambda[d_q(u_0, x_0) + d_q(Sx_0, Tu_0)] \\ &\leq (\beta + \lambda\gamma + \lambda\delta)d_q(x_1, u_1) + (\gamma\lambda + \delta\lambda)r. \end{aligned}$$

By using (3.10) and fact that  $u_0 \in \overline{B_{d_q}(x_0, r)}$ , in above inequality we obtain

$$\begin{aligned} d_q(x_2, u_2) &\leq (\beta + \lambda\gamma + \lambda\delta)\lambda r + (\gamma\lambda + \delta\lambda)r \\ &= (\beta + \lambda\gamma + \lambda\delta + \gamma + \delta)\lambda r = \lambda^2 r. \end{aligned}$$

Thus,

$$\begin{aligned} d_q(x_0, u_2) &\leq d_q(x_0, x_1) + d_q(x_1, x_2) + d_q(x_2, u_2) \\ &\leq d_q(x_0, x_1) + \lambda d_q(x_0, x_1) + \lambda^2 \leq r. \end{aligned}$$

Hence,  $u_2 \in \overline{B_{d_q}(x_0, r)}$ . Since  $T$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(x_0, Sx_0) = \alpha(u_2, u_3) \geq 1$ . Therefore, from  $(P'_2)$  holds. Suppose,  $(P'_1), (P'_2), \dots, (P'_i)$  be the inductive hypothesis. We shall show that  $(P'_{i+1})$  holds. For this, we consider two possible cases. First, suppose that  $i$  is even. Then, since  $\alpha(u_i, u_{i+1}) \geq 1$  and using (3.2), we get that

$$d_q(u_{i+1}, u_{i+2}) \leq \frac{\beta + \gamma + \delta}{1 - \gamma - \delta} d_q(u_i, u_{i+1}) = \lambda^{i+1} d_q(u_0, u_1).$$

Since  $\alpha(x_i, u_i) \geq 1$ , using (3.1) that

$$\begin{aligned} d_q(x_{i+1}, u_{i+1}) &= d_q(Sx_i, Tu_i) \\ &\leq \beta d_q(x_i, u_i) + \gamma [d_q(x_i, Sx_i) + d_q(u_i, Tu_i)] \\ &\quad + \delta [d_q(u_i, Sx_i) + d_q(x_i, Tu_i)] \\ &\leq \beta d_q(x_i, x_{i+1}) + \gamma \lambda [d_q(x_0, Sx_0) + d_q(u_0, Tu_0)] \\ &\quad + \delta \lambda [d_q(u_0, Sx_0) + d_q(x_0, Tu_0)] \end{aligned}$$

which gives with (c) and  $P'_i$

$$\begin{aligned} d_q(x_{i+1}, u_{i+1}) &\leq \beta d_q(x_i, u_i) + \gamma \lambda^i [d_q(x_0, u_0) + d_q(Sx_0, Tu_0)] \\ &\quad + \delta \lambda^i [d_q(u_0, x_0) + d_q(Sx_0, Tu_0)] \\ &\leq \beta \lambda^i r + \gamma \lambda^i [r + \lambda r] + \delta \lambda^i [r + \lambda r] \\ &= (\beta + \lambda \gamma + \lambda \delta + \gamma + \delta) \lambda^i r = \lambda^{i+1} r. \end{aligned}$$

Thus,

$$\begin{aligned} d_q(x_0, u_{i+1}) &\leq d_q(x_0, x_1) + d_q(x_1, x_2) + \dots + d_q(x_i, x_{i+1}) + d_q(x_{i+1}, u_{i+1}) \\ &\leq (1 + \lambda + \lambda^2 + \dots + \lambda^i) d_q(x_0, x_1) + \lambda^{i+1} r \\ &\leq (1 + \lambda + \lambda^2 + \dots + \lambda^i) (1 - \lambda) r + \lambda^{i+1} r = r. \end{aligned}$$

Hence,  $u_{i+1} \in \overline{B_{d_q}(x_0, r)}$ . Since  $S$  is an  $\alpha$ -dominated mapping on  $\overline{B_{d_q}(x_0, r)}$ , we have  $\alpha(u_{i+1}, Su_{i+1}) = \alpha(u_{i+1}, u_{i+2}) \geq 1$ . Also, since  $\alpha(x_{i+1}, u_0) \geq 1$ ,  $\alpha(u_n, u_{n+1}) \geq 1$ ,  $n = 0, 1, 2, \dots, i + 1$ , by triangular nature of  $\alpha$ , we have  $\alpha(x_{i+1}, u_{i+1}) \geq 1$ . Therefore,  $(P'_{i+1})$  holds. Similarly, one can see that if  $i$  is odd, then  $(P'_{i+1})$  holds,

which completes the inductive proof. Thus, for all  $n \in \mathbb{N}$ , we have

$$\begin{aligned}
 d_q(z, u_{2n}) &= d_q(Tz, Su_{2n-1}) \\
 &\leq \beta d_q(z, u_{2n-1}) + \gamma [d_q(z, Tz) + d_q(u_{2n-1}, Su_{2n-1})] \\
 &\quad + \delta [d_q(u_{2n-1}, Tz) + d_q(z, Su_{2n-1})] \\
 &\leq \beta d_q(z, u_{2n-1}) + \gamma d_q(u_{2n-1}, u_{2n}) \\
 &\quad + \delta [d_q(u_{2n-1}, z) + d_q(z, u_{2n})] \\
 &= (\beta + \delta) d_q(z, u_{2n-1}) + \gamma d_q(u_{2n-1}, u_{2n}) + \delta d_q(z, u_{2n}) \\
 &\leq (\beta + 2\delta) d_q(z, u_{2n-1}) + (\gamma + \delta) d_q(u_{2n-1}, u_{2n}) \\
 &\leq (\beta + 2\delta) d_q(Tz, Su_{2n-2}) + (\gamma + \delta) d_q(u_{2n-1}, u_{2n}) \\
 &\leq (\beta + 2\delta)^2 d_q(z, u_{2n-2}) + (\beta + 2\delta)(\gamma + \delta) d_q(u_{2n-2}, u_{2n-1}) \\
 &\quad + (\gamma + \delta) d_q(u_{2n-1}, u_{2n}) \\
 &\leq (\beta + 2\delta)^2 d_q(Tz, Su_{2n-3}) + (\beta + 2\delta)(\gamma + \delta) d_q(u_{2n-2}, u_{2n-1}) \\
 &\quad + (\gamma + \delta) d_q(u_{2n-1}, u_{2n}) \\
 &\leq (\beta + 2\delta)^3 d_q(z, u_{2n-3}) + (\beta + 2\delta)^2 (\gamma + \delta) d_q(u_{2n-3}, u_{2n-2}) \\
 &\quad + (\beta + 2\delta)(\gamma + \delta) d_q(u_{2n-2}, u_{2n-1}) + (\gamma + \delta) d_q(u_{2n-1}, u_{2n}) \\
 &\quad \vdots \\
 &\leq (\beta + 2\delta)^{2n} d_q(z, u_0) + (\beta + 2\delta)^{2n-1} (\gamma + \delta) d_q(u_0, u_1) + \dots \\
 &\quad + (\beta + 2\delta)(\gamma + \delta) d_q(u_{2n-2}, u_{2n-1}) + (\gamma + \delta) d_q(u_{2n-1}, u_{2n})
 \end{aligned}$$

Since  $\frac{\beta+2\delta}{\lambda} = \frac{(\beta+2\delta)(1-\gamma-\delta)}{\beta+\gamma+\delta} < 1$ , using  $(P'_n)$  in the above inequality we obtain

$$\begin{aligned}
 d_q(z, u_{2n}) &\leq (\beta + 2\delta)^{2n} d_q(z, u_0) + (\beta + 2\delta)^{2n-1} (\gamma + \delta) d_q(u_0, u_1) + \dots \\
 &\quad + (\beta + 2\delta)(\gamma + \delta) \lambda^{2n-2} d_q(u_0, u_1) + (\gamma + \delta) \lambda^{2n-1} d_q(u_0, u_1) \\
 &= (\beta + 2\delta)^{2n} d_q(z, u_0) \\
 &\quad + (\gamma + \delta) \lambda^{2n-1} d_q(u_0, u_1) \left[ 1 + \frac{\beta + 2\delta}{\lambda} + \dots + \left( \frac{\beta + 2\delta}{\lambda} \right)^{2n-1} \right] \\
 &\leq (\beta + 2\delta)^{2n} d_q(z, u_0) + \frac{(\gamma + \delta) \lambda^{2n-1} d_q(u_0, u_1)}{1 - \frac{\beta+2\delta}{\lambda}}
 \end{aligned}$$

Since  $\beta + 2\delta, \lambda \in [0, 1)$ , it follows from the above inequality that

$$\lim_{n \rightarrow \infty} d_q(z, u_{2n}) = 0. \quad (3.11)$$

Similarly, we can show that

$$\lim_{n \rightarrow \infty} d_q(u_{2n}, z) = \lim_{n \rightarrow \infty} d_q(u_{2n}, z^*) = \lim_{n \rightarrow \infty} d_q(z^*, u_{2n}) = 0. \quad (3.12)$$

By using (3.11) and (3.12), we obtain

$$d_q(z, z^*) \leq d_q(z, u_{2n}) + d_q(u_{2n}, z^*) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

$$d_q(z^*, z) \leq d_q(z^*, u_{2n}) + d_q(u_{2n}, z) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Hence,  $d_q(z, z^*) = d_q(z^*, z) = 0$ , i.e.,  $z = z^*$   $\square$

**Corollary 3.6.** *Let  $(X, d_q)$  be a left  $K$ -sequentially complete dislocated quasi metric space and  $T, S : X \rightarrow X$  be two mappings. Let  $x_0 \in X, r > 0, x_0 \in \overline{B_{d_q}(x_0, r)}$  and*

there exist nonnegative real numbers  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + 2\delta \in (0, 1)$  and the following conditions hold:

$$d_q(Sx, Ty) \leq \beta d_q(x, y) + \gamma[d_q(x, Sx) + d_q(y, Ty)] + \delta[d_q(y, Sx) + d_q(x, Ty)],$$

$$d_q(Tx, Sy) \leq \beta d_q(x, y) + \gamma[d_q(x, Tx) + d_q(y, Sy)] + \delta[d_q(y, Tx) + d_q(x, Sy)],$$

and

$$d_q(x_0, Sx_0) \leq (1 - \lambda)r,$$

where  $\lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta}$ . Then there exists a unique point  $z \in \overline{B_{d_q}(x_0, r)}$  such that  $z = Sz = Tz$  and  $d_q(z, z) = 0$ . Moreover,  $S$  and  $T$  have no fixed point in  $\overline{B_{d_q}(x_0, r)}$  other than  $z$ .

*Proof.* The proof follows by the previous results, taking  $\alpha : X \times X \rightarrow [0, \infty)$  with  $\alpha(x, y) = 1$  for all  $x, y \in X$ .  $\square$

**Theorem 3.7.** Let  $(X, d_q)$  be a left  $K$ -sequentially complete dislocated quasi metric space. Suppose, there exist a function  $\alpha : X \times X \rightarrow [0, +\infty)$  and nonnegative constants  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + 2\delta \in (0, 1)$  and the following conditions hold:

$$d_q(Sx, Ty) \leq \beta d_q(x, y) + \gamma[d_q(x, Sx) + d_q(y, Ty)] + \delta[d_q(y, Sx) + d_q(x, Ty)],$$

$$d_q(Tx, Sy) \leq \beta d_q(x, y) + \gamma[d_q(x, Tx) + d_q(y, Sy)] + \delta[d_q(y, Tx) + d_q(x, Sy)],$$

for all  $x, y \in X$  such that  $\alpha(x, y) \geq 1$  or  $\alpha(y, x) \geq 1$ . If  $(X, d_q)$  is  $\alpha$ -regular, then there exists a point  $z$  in  $X$  such that  $z = Sz = Tz$  and  $d_q(z, z) = 0$ .

*Proof.* By Theorem 3.1, the condition (3.3) is imposed in order to restrict the contractive conditions (3.1) and (3.2) to  $\overline{B_{d_q}(x_0, r)}$ . However, the condition (3.3) can be relaxed by imposing the conditions (3.1) and (3.2) to all elements  $x, y \in X$  such that  $\alpha(x, y) \geq 1$  or  $\alpha(y, x) \geq 1$ , we obtain the following result.  $\square$

Recall that, if  $(X, \preceq)$  is a pre-ordered set and  $T : X \rightarrow X$  is such that  $Tx = x$  for all  $x \in A \subseteq X$ , then the mapping  $T$  is said to be dominated on  $A$ . Define the set  $\nabla$  by

$$\nabla = \{(x, y) \in X \times X : x \preceq y \text{ or } y \preceq x\}.$$

**Theorem 3.8.** Let  $(X, \preceq, d_q)$  be a pre-ordered left  $K$ -sequentially complete dislocated quasi metric space,  $x_0 \in X$ ,  $r > 0$  and  $S, T : X \rightarrow X$  be two dominated mappings on  $\overline{B_{d_q}(x_0, r)}$ . Suppose that there exist nonnegative real numbers  $\beta, \gamma, \delta$  such that  $\beta + 2\gamma + \delta \in (0, 1)$  and the following conditions hold:

$$d_q(Sx, Ty) \leq \beta d_q(x, y) + \gamma[d_q(x, Sx) + d_q(y, Ty)] + \delta[d_q(y, Sx) + d_q(x, Ty)],$$

$$d_q(Tx, Sy) \leq \beta d_q(x, y) + \gamma[d_q(x, Tx) + d_q(y, Sy)] + \delta[d_q(y, Tx) + d_q(x, Sy)],$$

for all  $(x, y) \in \overline{B_{d_q}(x_0, r)} \times \overline{B_{d_q}(x_0, r)} \cap \nabla$  and

$$d_q(x_0, Sx_0) \leq (1 - \lambda)r,$$

where  $\lambda = \frac{\beta + \gamma + \delta}{1 - \gamma - \delta}$ . If for any sequence  $\{x_n\} \in \overline{B_{d_q}(x_0, r)}$  such that  $(x_n, x_{n+1}) \in \nabla$ ,  $x_n \rightarrow w$  as  $n \rightarrow \infty$  implies that  $(w, x_n) \in \nabla$  for all  $n \geq 0$ , then there exists a point  $z \in \overline{B_{d_q}(x_0, r)}$  such that  $z = Sz = Tz$  and  $d_q(z, z) = 0$ . In addition, suppose that:

(a)  $(x, y), (y, z) \in \nabla$  implies  $(x, z) \in \nabla$ .

(b) For  $x, y \in \overline{B_{d_q}(x_0, r)}$  there exists  $u_0 \in \overline{B_{d_q}(x_0, r)}$  such that  $(x, u_0), (y, u_0) \in \nabla$ .

(c) For all  $u \in \overline{B_{d_q}(x_0, r)}$  such that  $(u, Sx_0) \in \nabla$  the following condition holds  $d_q(x_0, Sx_0) + d_q(u, Tu) + d_q(u, Sx_0) + d_q(x_0, Tu) \leq d_q(x_0, u) + d_q(Sx_0, Tu)$ .

Then,  $z$  is the unique common fixed point of  $S$  and  $T$  in  $\overline{B_{d_q}(x_0, r)}$ .

*Proof.* This follows from Theorem 3.6 taking  $\alpha : X \times X \rightarrow [0, +\infty)$  defined as

$$\alpha(x, y) = \begin{cases} 1, & \text{If } (x, y) \in \nabla, \\ 0, & \text{otherwise.} \end{cases}$$

□

#### 4. CONCLUSIONS

We prov some common fixed point theorems for mappings under Hardy Rogers contractive conditions on a left  $K$ -sequentially complete dislocated quasi metric space.

#### ACKNOWLEDGMENTS

The first and third authors thank for the support of Rambhai Barni Rajabhat University. Finally, Dhananjay Gopal was supported by Sardar Vallabhbhai National Institute of Technology.

#### REFERENCES

1. A Amini-Harandi, Metric-likespaces, partial metric spaces and fixed points, Fixed Point Theory Appl. **2012**, 2012:204.
2. M. Arshad, A. Azam, M. Abbas and A. Shoaib, Fixed point results of dominated mappings on a closed ball in ordered partial metric spaces without continuity, U.P.B. Sci. Bull., Series A. **76** (2) (2014).
3. M. Arshad, Z. Kadelburg, S. Radenović, A. Shoaib, S. Shukla, Fixed Points of  $\alpha$ -Dominated Mappings on Dislocated Quasi Metric Spaces, Filomat **31:11** (2017), 3041-3056.
4. M. Arshad, A. Shoaib and P. Vetro, Common fixed points of a pair Of Hardy Rogers type mappings on a closed ball in ordered dislocated metric spaces, J. Funct. Spaces Appl. **2013** Article ID 63818 (2013).
5. M. Arshad, A. Shoaib, M. Abbas and A. Azam, Fixed points of a pair of Kannan type mappings on a closed ball in ordered partial metric spaces, Misk. Math. Notes, **14** (2013), 769–784.
6. H. Aydi, Some fixed point results in ordered partial metric spaces, J. Nonlinear Sci. Appl. **4** (2011), 210-217.
7. I. Beg, M. Arshad and A. Shoaib, Fixed point in a closed ball in ordered dislocated quasi metric space, Fixed Point Theory, **16**(2) (2015), 195-206 .
8. L.J.B. Ćirić A generalization of Banach's contraction principle, Proc. Amer. Math. Soc. **45** (1974), 267-273.
9. L. Ćirić, M. Abbas, R. Saadati and N. Hussain, Common fixed points of almost generalized contractive mappings in ordered metric spaces, Appl. Math. Comput. **217** (2011), 5784-5789.
10. E. Karapinar, İ. M. Erhan and A. Öztürk, Fixed point theorems on quasi-partial metric spaces, Math. Comp. Model. **57**(9-10) (2013), 2442-2448.
11. I.L. Reilly, P.V. Semirahmanya and M.K. Vamanamurthy, Cauchy sequences in quasi-pseudometric spaces, Monatsh. Math. **93** (1982), 127-140.
12. A. Shoaib, M. Arshad and J. Ahmad, Fixed point results of locally cotractive mappings in ordered quasi-partial metric spaces, Sci. World J. **2013** Article ID 194897, 8 pages (2013).
13. F.M. Zeyada, G.H. Hassan and M.A. Ahmed, A generalization of a fixed point theorem due to Hitzler and Seda in dislocated quasi-metric spaces, Arabian J. Sci. Eng. A. **31**(1) (2006), 111–114.
14. P. Hitzler, Generalized Metrics and Topology in Logic Programming Semantics, Ph.D. Thesis, National University of Ireland (University College, Cork), 2001.
15. N. Hussain M. Arshad, A. Shoaib and Fahimuddin, Common fixed point results for  $\alpha$ - $\psi$  contractions on a metric space endowed with graph, J. Inequ. Appl., **(2014)** 2014:136.
16. N. Hussain, J. R. Roshan, V. Parvaneh and M. Abbas, Common fixed point results for weak contractive mappings in ordered b-dislocated metric spaces with applications, J. Inequ. Appl. **(2013)** 2013:486.

17. S. K. Malhotra, S. Radenović and S. Shukla, Some fixed point results without monotone property in partially ordered metric-like spaces, *Egyptian Math. Soc.*, **22** (2014) 83-89.
18. I.R. Sarma and P.S. Kumari, On dislocated metric spaces, *Int. J. Math. Arch.*, **3**(1) (2012) 72-77.
19. B. Samet, C. Vetro and P. Vetro, Fixed point theorems for  $\alpha$ - $\psi$ -contractive type mappings, *Nonlinear Anal.*, **75** (2012) 2154-2165.